


THRSP Goat anti-Human Polyclonal Antibody - LS-B3813 - LSBio


CatalogID:	LS-B3813
Validation:	This antibody replaces catalog number LS-C108168. It has been validated for use in the following assays: IHC-P.
Target:	thyroid hormone responsive (THRSP)
Synonyms:	THRSP Antibody, Lpgp Antibody, Lipogenic protein 1 Antibody, S14 Antibody, SPOT14 Antibody, THRP Antibody, SPOT14 homolog Antibody, LPGP1 Antibody, Spot 14 protein Antibody, Thyroid hormone responsive Antibody
Host	THRSP antibody was produced in Goat
Clonality:	Polyclonal
Immunogen Species:	THRSP antibody was raised against Human
Immunogen:	THRSP antibody was raised against synthetic peptide from human THRSP. Percent identity with other species by BLAST analysis: Human (100%), Gorilla (93%), Monkey (86%).
Specificity:	Human THRSP.
Reactivity:	Human
Predicted Reactivity:	Gorilla
Purification:	Immunoaffinity purified
Presentation:	Tris-buffered saline, pH 7.3, 0.5% BSA, 0.02% sodium azide
Recommended Storage:	Store at -20°C. Minimize freezing and thawing.
Usage Summary:	Immunohistochemistry: LS-B3813 was validated for use in immunohistochemistry on a panel of 21 formalin-fixed, paraffin-embedded (FFPE) human tissues after heat induced antigen retrieval in pH 6.0 citrate buffer. After incubation with the primary antibody, slides were incubated with biotinylated secondary antibody, followed by alkaline phosphatase-streptavidin and chromogen. The stained slides were evaluated by a pathologist to confirm staining specificity. The optimal working concentration for LS-B3813 was determined to be 3.75ug/ml.
Uses:	IHC - Paraffin (3.75 µg/ml) (Optimal dilution to be determined by the researcher)
Size:	50 µg
Concentration:	0.5 mg/ml

Immunohistochemistry Image:



Anti-THRSP antibody IHC of human thyroid. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B3813 concentration 3.75 ug/ml.

Immunohistochemistry Image:



Anti-THRSP antibody IHC of human breast. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B3813 concentration 3.75 ug/ml.

Requested From:

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/23/2014

© 2014 LifeSpan BioSciences