



NCKAP1L Goat anti-Human Polyclonal Antibody - LS-B3817 - LSBio	
CatalogID:	LS-B3817
Validation:	This antibody replaces catalog number LS-C108187. It has been validated for use in the following assays: IHC-P.
Target:	NCK-associated protein 1-like (NCKAP1L)
Synonyms:	NCKAP1L Antibody, HEM1 Antibody, Hematopoietic protein 1 Antibody, NCK-associated protein 1-like Antibody
Host	NCKAP1L antibody was produced in Goat
Clonality:	Polyclonal
Immunogen Species:	NCKAP1L antibody was raised against Human
Immunogen:	NCKAP1L antibody was raised against synthetic peptide from human NCKAP1L. Percent identity by BLAST analysis: Human, Gorilla, Gibbon, Monkey, Marmoset, Mouse, Rat, Hamster, Panda, Bovine, Dog, Bat, Horse, Rabbit (100%); Elephant (92%); Xenopus (85%).
Specificity:	Human NCKAP1L.
Reactivity:	Human, Gorilla, Gibbon, Monkey, Mouse, Rat, Bat, Bovine, Dog, Hamster, Horse, Rabbit
Purification:	Immunoaffinity purified
Presentation:	Tris-buffered saline, pH 7.3, 0.5% BSA, 0.02% sodium azide
Recommended Storage:	Store at -20°C. Minimize freezing and thawing.
Usage Summary:	Immunohistochemistry: LS-B3817 was validated for use in immunohistochemistry on a panel of 21 formalin-fixed, paraffin-embedded (FFPE) human tissues after heat induced antigen retrieval in pH 6.0 citrate buffer. After incubation with the primary antibody, slides were incubated with biotinylated secondary antibody, followed by alkaline phosphatase-streptavidin and chromogen. The stained slides were evaluated by a pathologist to confirm staining specificity. The optimal working concentration for LS-B3817 was determined to be 3.75ug/ml.
Uses:	IHC - Paraffin (3.75 µg/ml) (Optimal dilution to be determined by the researcher)
Size:	50 µg
Concentration:	0.5 mg/ml

Immunohistochemistry Image:



Anti-NCKAP1L antibody IHC of human spleen. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B3817 concentration 3.75 ug/ml.

Immunohistochemistry Image:



Anti-NCKAP1L antibody IHC of human colon. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B3817 concentration 3.75 ug/ml.

Requested From:

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/23/2014

© 2014 LifeSpan BioSciences