



NCOA3 / SRC-3 / AIB1 Goat anti-Human Polyclonal Antibody - LS-B3824 - LSBio	
CatalogID:	LS-B3824
Validation:	This antibody replaces catalog number LS-C108213. It has been validated for use in the following assays: IHC-Fr.
Target:	nuclear receptor coactivator 3 (NCOA3)
Synonyms:	NCOA3 Antibody, ACTR Antibody, AIB1 Antibody, BHLHe42 Antibody, CAGH16 Antibody, CBP-interacting protein Antibody, CTG26 Antibody, AIB-1 Antibody, P/CIP Antibody, TNRC14 Antibody, TRAM-1 Antibody, Nuclear receptor coactivator 3 Antibody, PCIP Antibody, RAC-3 Antibody, SRC3 Antibody, TNRC16 Antibody, KAT13B Antibody, NCoA-3 Antibody, SRC-3 Antibody
Host	NCOA3 antibody was produced in Goat
Clonality:	Polyclonal
Immunogen Species:	NCOA3 / SRC-3 / AIB1 antibody was raised against Human
Antigen Type:	Synthetic peptide
Immunogen:	NCOA3 / SRC-3 / AIB1 antibody was raised against synthetic peptide C-TNETQRQKSHT from an internal region of human NCOA3 (NP_006525.2; NP_858045.1). Percent identity by BLAST analysis: Human, Gorilla, Gibbon, Monkey, Marmoset, Hamster, Elephant, Panda, Dog, Bovine, Bat, Pig (100%); Mouse, Rat, Horse, Rabbit (91%); Opossum, Turkey, Platypus, Lizard (82%).
Specificity:	Human NCOA3. This antibody is expected to recognise both reported isoforms (NP_006525.2; NP_858045.1)
Reactivity:	Human, Gorilla, Gibbon, Monkey, Bat, Bovine, Dog, Hamster, Pig
Predicted Reactivity:	Mouse, Rat, Horse, Rabbit
Purification:	Immunoaffinity purified
Presentation:	Tris-buffered saline, pH 7.3, 0.5% BSA, 0.02% sodium azide
Recommended Storage:	Store at -20°C. Minimize freezing and thawing.
Usage Summary:	Immunohistochemistry: LS-B3824 was validated for use in immunohistochemistry on a panel of 21 formalin-fixed, paraffin-embedded (FFPE) human tissues after heat induced antigen retrieval in pH 6.0 citrate buffer. After incubation with the primary antibody, slides were incubated with biotinylated secondary antibody, followed by alkaline phosphatase-streptavidin and chromogen. The stained slides were evaluated by a pathologist to confirm staining specificity. The optimal working concentration for LS-B3824 was determined to be 3.75ug/ml.
Uses:	IHC - Paraffin (3.75 µg/ml) (Optimal dilution to be determined by the researcher)
Size:	50 µg
Concentration:	0.5 mg/ml

Immunohistochemistry Image:



Anti-NCOA3 antibody IHC of human spleen. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B3824 concentration 3.75 ug/ml.

Immunohistochemistry Image:



Anti-NCOA3 antibody IHC of human placenta. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B3824 concentration 3.75 ug/ml.

Requested From:

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/23/2014

© 2014 LifeSpan BioSciences