


TRIB2 Goat anti-Human Polyclonal Antibody - LS-B3826 - LSBio


CatalogID:	LS-B3826
Validation:	This antibody replaces catalog number LS-C108224. It has been validated for use in the following assays: IHC-P.
Target:	tribbles pseudokinase 2 (TRIB2)
Synonyms:	TRIB2 Antibody, C5FW Antibody, GS3955 Antibody, TRB-2 Antibody, TRB2 Antibody, Tribbles homolog 2 Antibody
Host	TRIB2 antibody was produced in Goat
Clonality:	Polyclonal
Immunogen Species:	TRIB2 antibody was raised against Human
Immunogen:	TRIB2 antibody was raised against synthetic peptide from human TRIB2. Percent identity by BLAST analysis: Human, Gorilla, Orangutan, Gibbon, Marmoset, Mouse, Rat, Elephant, Panda, Bat, Bovine, Dog, Horse, Rabbit, Pig, Xenopus (100%); Turkey, Chicken, Platypus (92%); Opossum (85%).
Specificity:	Human TRIB2.
Reactivity:	Human, Gorilla, Orangutan, Gibbon, Monkey, Mouse, Rat, Bat, Bovine, Dog, Horse, Pig, Rabbit, Xenopus
Predicted Reactivity:	Chicken
Purification:	Immunoaffinity purified
Presentation:	Tris-buffered saline, pH 7.3, 0.5% BSA, 0.02% sodium azide
Recommended Storage:	Store at -20°C. Minimize freezing and thawing.
Usage Summary:	Immunohistochemistry: LS-B3826 was validated for use in immunohistochemistry on a panel of 21 formalin-fixed, paraffin-embedded (FFPE) human tissues after heat induced antigen retrieval in pH 6.0 citrate buffer. After incubation with the primary antibody, slides were incubated with biotinylated secondary antibody, followed by alkaline phosphatase-streptavidin and chromogen. The stained slides were evaluated by a pathologist to confirm staining specificity. The optimal working concentration for LS-B3826 was determined to be 3.75ug/ml.
Uses:	IHC - Paraffin (3.75 µg/ml) (Optimal dilution to be determined by the researcher)
Size:	50 µg
Concentration:	0.5 mg/ml

Immunohistochemistry Image:



Anti-TRIB2 antibody IHC of human spleen. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B3826 concentration 3.75 ug/ml.

Immunohistochemistry Image:



Anti-TRIB2 antibody IHC of human colon. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B3826 concentration 3.75 ug/ml.

Requested From:

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/23/2014

© 2014 LifeSpan BioSciences