


| CUL9 / Cullin 9 Rabbit anti-Human Polyclonal (C-Terminus) Antibody - LS-B401 - LSBio |   |
|--|---|
|  |   |
| CatalogID:   | LS-B401   |
| Validation:  | This antibody replaces catalog number LS-C19122. It has been validated for use in the following assays: IHC.  |
| Target:  | cullin 9 (CUL9)   |
| Synonyms:  | CUL9 Antibody, Cullin 9 Antibody, Cullin-9 Antibody, CUL-9 Antibody, H7AP1 Antibody, RP3-330M21.2 Antibody, KIAA0708 Antibody, PARC Antibody, UbcH7-associated protein 1 Antibody   |
| Host   | CUL9 antibody was produced in Rabbit  |
| Clonality:   | Polyclonal  |
| Immunogen Species:   | CUL9 / Cullin 9 antibody was raised against Human   |
| Antigen Type:  | Synthetic peptide   |
| Immunogen:   | CUL9 / Cullin 9 antibody was raised against synthetic peptide from human CUL9 / Cullin 9.   |
| Specificity:   | Amino acids 2503-2517 of Human PARC (C-terminus) coupled to KLH.  |
| Epitope:   | C-Terminus  |
| Reactivity:  | Human   |
| Purification:  | Delipidated and defibrinated  |
| Presentation:  | Sterile filtered antiserum, 0.01% sodium azide.   |
| Recommended Storage:   | +4°C or -20°C, Avoid repeated freezing and thawing.   |
| Usage Summary:   | Immunohistochemistry: LS-B401 was validated for use in immunohistochemistry on a panel of 21 formalin-fixed, paraffin-embedded (FFPE) human tissues after heat induced antigen retrieval in pH 6.0 citrate buffer. After incubation with the primary antibody, slides were incubated with biotinylated secondary antibody, followed by alkaline phosphatase-streptavidin and chromogen. The stained slides were evaluated by a pathologist to confirm staining specificity. The optimal working concentration for LS-B401 was determined to be 1:500. |
| Uses:  | IHC - Paraffin (1:500), Western blot (1:500 - 1:1000), Immunoprecipitation, ELISA (1:2000 - 1:10000) (Optimal dilution to be determined by the researcher)  |
| Size:  | 50 μl   |

## Immunohistochemistry Image:



Anti-CUL9 / Cullin 9 antibody IHC of human colon. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B401 dilution 1:500.

Requested From: Japan

Laboratory Reagent For In Vitro Research Use Only
Not for resale without prior written consent from LifeSpan BioSciences, Inc.
Created on 9/23/2014
© 2014 LifeSpan BioSciences