



BAK1 / BAK Rabbit anti-Human Polyclonal Antibody - LS-B4072 - LSBio	
CatalogID:	LS-B4072
Validation:	This antibody replaces catalog number LS-C109284. It has been validated for use in the following assays: IHC-P.
Target:	BCL2-antagonist/killer 1 (BAK1)
Synonyms:	BAK1 Antibody, Apoptosis regulator BAK Antibody, BAK Antibody, BCL2L7 Antibody, Bcl-2-like protein 7 Antibody, BCL2-antagonist/killer 1 Antibody, Bcl2-L-7 Antibody, BCL2-like 7 protein Antibody, CDN1 Antibody, Pro-apoptotic protein BAK Antibody, BAK-LIKE Antibody
Host	BAK1 antibody was produced in Rabbit
Clonality:	Polyclonal
Immunogen Species:	BAK1 / BAK antibody was raised against Human
Antigen Type:	Synthetic peptide
Immunogen:	BAK1 / BAK antibody was raised against synthetic peptide containing a sequence corresponding to a region within amino acids 1 and 14 of Human BAK1.
Specificity:	Human BAK1.
Reactivity:	Human
Purification:	Immunoaffinity purified
Presentation:	PBS, pH 7, 1% BSA, 20% Glycerol, 0.01% Thimerosal
Recommended Storage:	Aliquot and store at -20°C. Minimize freezing and thawing.
Uses:	IHC - Paraffin (10 µg/ml), Western blot (1:500 - 1:3000) (Optimal dilution to be determined by the researcher)
Size:	50 µl

Immunohistochemistry Image:




Anti-BAK1 / BAK antibody IHC of human liver, bile duct. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B4072 concentration 10 ug/ml.

Immunohistochemistry Image:




Anti-BAK1 / BAK antibody IHC of human lung, respiratory epithelium. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B4072 concentration 10 ug/ml.

Immunohistochemistry Image:



Anti-BAK1 / BAK antibody IHC of human breast. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B4072 concentration 10 ug/ml.

Western Blot Image:



Sample (30 ug of whole cell lysate). A: HeLa. B: Hep G2. C: Molt-4. 12% SDS PAGE. BAK1 antibody diluted at 1:1000.

Requested From:	Japan
Laboratory Reagent For In Vitro Research Use Only	
Not for resale without prior written consent from LifeSpan BioSciences, Inc.	
Created on 9/23/2014	
© 2014 LifeSpan BioSciences	