



HTR2C / 5-HT2C Receptor Rabbit anti-Human Polyclonal Antibody - LS-B4082 - LSBio	
<b>CatalogID:</b>	LS-B4082
<b>Validation:</b>	This antibody replaces catalog number LS-C109302. It has been validated for use in the following assays: IHC-P.
<b>Target:</b>	5-hydroxytryptamine (serotonin) receptor 2C, G protein-coupled (HTR2C)
<b>Synonyms:</b>	HTR2C Antibody, 5-HT-1C Antibody, 5-HT1c receptor Antibody, 5HT-1C Antibody, 5-HT2C Antibody, 5-HT2c receptor Antibody, 5HT2C Receptor Antibody, 5-HT-2C Antibody, 5-HT1C Antibody, 5-HTR2C Antibody, HTR1C Antibody, Serotonin 5-HT-1c receptor Antibody, Serotonin receptor 2C Antibody, Serotonin 1c receptor Antibody, Serotonin 5-HT-2C receptor Antibody, Serotonin 2c receptor Antibody
<b>Family / Subfamily:</b>	GPCR / Serotonin
<b>Host</b>	HTR2C antibody was produced in Rabbit
<b>Clonality:</b>	Polyclonal
<b>Immunogen Species:</b>	HTR2C / 5-HT2C Receptor antibody was raised against Human
<b>Antigen Type:</b>	Synthetic peptide
<b>Immunogen:</b>	HTR2C / 5-HT2C Receptor antibody was raised against synthetic peptide containing a sequence corresponding to a region within amino acids 387 and 446 of Human 5HT2C Receptor.
<b>Specificity:</b>	Human 5HT2C receptor.
<b>Reactivity:</b>	Human
<b>Purification:</b>	Immunoaffinity purified
<b>Presentation:</b>	0.1 M Tris-glycine, pH 7, 10% Glycerol, 0.01% Thimerosal
<b>Recommended Storage:</b>	Aliquot and store at -20°C. Minimize freezing and thawing.
<b>Uses:</b>	IHC - Paraffin (10 µg/ml), Western blot (1:500 - 1:3000) (Optimal dilution to be determined by the researcher)
<b>Size:</b>	50 µl

**Immunohistochemistry Image:**



Anti-5HT2C Receptor antibody IHC of human prostate. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B4082 concentration 10 ug/ml.

**Immunohistochemistry Image:**



Anti-5HT<sub>2C</sub> Receptor antibody IHC of human skeletal muscle. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B4082 concentration 10 ug/ml.

**Requested From:**

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/23/2014

© 2014 LifeSpan BioSciences