


TF / Transferrin Rabbit anti-Human Polyclonal Antibody - LS-B426 - LSBio

CatalogID:	LS-B426
Validation:	This antibody replaces catalog number LS-C18894. It has been validated for use in the following assays: IHC.
Target:	transferrin (TF)
Synonyms:	TF Antibody, PRO1557 Antibody, Serotransferrin Antibody, Siderophilin Antibody, TFQTL1 Antibody, Beta-1 metal-binding globulin Antibody, Transferrin Antibody, PRO2086 Antibody
Host	TF antibody was produced in Rabbit
Clonality:	Polyclonal
Immunogen Species:	TF / Transferrin antibody was raised against Human
Specificity:	Human Transferrin.
Reactivity:	Human
Purification:	Immunoaffinity purified
Presentation:	0.02 M potassium phosphate, 0.15 M sodium chloride, pH 7.2, 0.01% sodium azide.
Recommended Storage:	+4°C or -20°C, Avoid repeated freezing and thawing.
Usage Summary:	Immunohistochemistry: LS-B426 was validated for use in immunohistochemistry on a panel of 21 formalin-fixed, paraffin-embedded (FFPE) human tissues after heat induced antigen retrieval in pH 6.0 citrate buffer. After incubation with the primary antibody, slides were incubated with biotinylated secondary antibody, followed by alkaline phosphatase-streptavidin and chromogen. The stained slides were evaluated by a pathologist to confirm staining specificity. The optimal working concentration for LS-B426 was determined to be 2.5 ug/ml.
Uses:	IHC - Paraffin (2.5 µg/ml), Immunofluorescence, ELISA (Optimal dilution to be determined by the researcher)
Size:	50 µg

Immunohistochemistry Image:



Anti-TF / Transferrin antibody IHC of human brain, cortex. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B426 concentration 5 ug/ml.

Requested From:

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/23/2014

© 2014 LifeSpan BioSciences