



ANAPC13 Rabbit anti-Human Polyclonal (Internal) Antibody - LS-B4648 - LSBio	
CatalogID:	LS-B4648
Validation:	This antibody replaces catalog number LS-C108715. It has been validated for use in the following assays: IHC-P.
Target:	anaphase promoting complex subunit 13 (ANAPC13)
Synonyms:	ANAPC13 Antibody, APC13 Antibody, Cylosome subunit 13 Antibody, Cyclosome subunit 13 Antibody, SWM1 Antibody, DKFZP566D193 Antibody
Host	ANAPC13 antibody was produced in Rabbit
Clonality:	Polyclonal
Immunogen Species:	ANAPC13 antibody was raised against Human
Antigen Type:	Synthetic peptide
Immunogen:	ANAPC13 antibody was raised against a 17 amino acid peptide near the center of human APC13.
Specificity:	Human ANAPC13
Epitope:	Internal
Reactivity:	Human, Mouse, Rat
Purification:	Immunoaffinity purified
Presentation:	PBS, 0.02% sodium azide
Recommended Storage:	Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles. Store undiluted.
Usage Summary:	APC13 antibody can be used for detection of APC13 by Western blot at 1-2 ug/ml.
Uses:	IHC - Paraffin (5 μg/ml), Western blot (1 - 2 μg/ml), ELISA (Optimal dilution to be determined by the researcher)
Size:	50 μg
Concentration:	1 mg/ml

Immunohistochemistry Image:




Anti-ANAPC13 antibody IHC of human adrenal. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B4648 concentration 5 ug/ml.

Immunofluorescence Image:



Immunofluorescence of APC13 in Jurkat cells with APC13 antibody at 20 ug/ml.

Western Blot Image:



Western blot of APC13 in Jurkat cell tissue lysate with APC13 antibody at 1 ug/ml in (A) the absence and (B) the presence of blocking peptide.

Requested From:

Japan

Laboratory Reagent For In Vitro Research Use Only
Not for resale without prior written consent from LifeSpan BioSciences, Inc.
Created on 9/23/2014
© 2014 LifeSpan BioSciences