



FNBP1 / FBP17 Goat anti-Human Polyclonal (Internal) Antibody - LS-B4929 - LSBio	
CatalogID:	LS-B4929
Validation:	This antibody replaces catalog number LS-C54820. It has been validated for use in the following assays: IHC-P.
Target:	formin binding protein 1 (FNBP1)
Synonyms:	FNBP1 Antibody, FBP17 Antibody, HFBP17 Antibody, KIAA0554 Antibody, Formin-binding protein 1 Antibody, Formin-binding protein 17 Antibody, Formin binding protein 1 Antibody
Host	FNBP1 antibody was produced in Goat
Clonality:	Polyclonal
Immunogen Species:	FNBP1 / FBP17 antibody was raised against Human
Antigen Type:	Synthetic peptide
Immunogen:	FNBP1 / FBP17 antibody was raised against synthetic peptide KQLESSKRRFERDC from an internal region of human FNBP1 (NP_055848.1). Percent identity by BLAST analysis: Human, Gibbon, Monkey, Marmoset, Mouse, Rat, Hamster, Elephant, Panda, Dog, Bat, Bovine, Horse, Pig, Opossum, Zebrafish (100%); Turkey, Chicken, Platypus, Xenopus, Pufferfish (93%); Gorilla, Orangutan (86%).
Specificity:	Human FNBP1.
Epitope:	Internal
Reactivity:	Human, Gibbon, Monkey, Mouse, Rat, Bat, Bovine, Dog, Hamster, Horse, Pig, Zebrafish
Predicted Reactivity:	Chicken, Xenopus
Purification:	Immunoaffinity purified
Presentation:	Tris-buffered saline, pH 7.3, 0.5% BSA, 0.02% sodium azide
Recommended Storage:	Store at -20°C. Minimize freezing and thawing.
Uses:	IHC - Paraffin (3.75 µg/ml), Western blot (1:32000) & (0.25 - 2 µg/ml), ELISA (1:32000) (Optimal dilution to be determined by the researcher)
Size:	50 µg
Concentration:	0.5 mg/ml

Immunohistochemistry Image:




Anti-FBNP1 antibody IHC of human small intestine. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B4929 concentration 3.75 ug/ml.

Immunohistochemistry Image:



Anti-FNBP1 antibody IHC of human tonsil. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B4929 concentration 3.75 ug/ml.

Western Blot Image:



Antibody staining (0.25 ug/ml) of human testis lysate (RIPA buffer, 35 ug total protein per lane). Primary incubated for 1 hour. Detected by Western blot of chemiluminescence.

Requested From:

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/24/2014

© 2014 LifeSpan BioSciences