



VPS37C Goat anti-Human Polyclonal (Internal) Antibody - LS-B4930 - LSBio	
CatalogID:	LS-B4930
Validation:	This antibody replaces catalog number LS-C55678. It has been validated for use in the following assays: IHC-P.
Target:	vacuolar protein sorting 37 homolog C (S. cerevisiae) (VPS37C)
Synonyms:	VPS37C Antibody, HVps37C Antibody, PML39 Antibody, Vacuolar protein sorting 37C Antibody, ESCRT-I complex subunit VPS37C Antibody
Host	VPS37C antibody was produced in Goat
Clonality:	Polyclonal
Immunogen Species:	VPS37C antibody was raised against Human
Antigen Type:	Synthetic peptide
Immunogen:	VPS37C antibody was raised against synthetic peptide C-ETLKDKTLQELEELQ from an internal region of human VPS37C (NP_060436.4). Percent identity by BLAST analysis: Human, Gorilla, Orangutan, Gibbon, Marmoset (100%); Monkey, Bat, Rabbit (93%); Mouse, Rat, Hamster, Bovine, Dog, Horse (87%); Elephant (80%).
Specificity:	Human VPS37C.
Epitope:	Internal
Reactivity:	Human, Gorilla, Orangutan, Gibbon
Predicted Reactivity:	Monkey, Bat, Rabbit
Purification:	Immunoaffinity purified
Presentation:	Tris-buffered saline, pH 7.3, 0.5% BSA, 0.02% sodium azide
Recommended Storage:	Store at -20°C. Minimize freezing and thawing.
Uses:	IHC - Paraffin (3.75 μg/ml), Western blot (1:32000) & (0.1 μg/ml), ELISA (1:32000) (Optimal dilution to be determined by the researcher)
Size:	50 µg
Concentration:	0.5 mg/ml

Immunohistochemistry Image:



Anti-VPS37C antibody IHC of human prostate. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B4930 concentration 3.75 ug/ml.

Immunohistochemistry Image:



Anti-VPS37C antibody IHC of human breast. Immunohistochemistry of formalin-fixed, paraffin -embedded tissue after heat-induced antigen retrieval. Antibody LS-B4930 concentration 3.75 ug/ml.

Western Blot Image:		
250kDa 150kDa 100kDa		
75kDa 50kDa		
37kDa		
25kDa		
20kDa		
15kDa		
10kDa		
Antibody (0.03g/ml) sta Primary incubation was	aining of human duodenum lysate (35 ug protein in RIPA buffer). s 1 hour. Detected by chemiluminescence.	
Requested From:	Japan	
Labor	atory Reagent For In Vitro Research Use Only	
Not for resale without prior written consent from LifeSpan BioSciences, Inc.		
Created on 9/24/2014		
	© 2014 LifeSpan BioSciences	