


TRIM7 Goat anti-Human Polyclonal (C-Terminus) Antibody - LS-B4968 - LSBio


CatalogID:	LS-B4968
Validation:	This antibody replaces catalog number LS-C54922. It has been validated for use in the following assays: IHC-P.
Target:	tripartite motif containing 7 (TRIM7)
Synonyms:	TRIM7 Antibody, Glycogenin-interacting protein Antibody, GNIP Antibody, RING finger protein 90 Antibody, RNF90 Antibody, Tripartite motif protein TRIM7 Antibody, Tripartite motif-containing 7 Antibody, Tripartite motif containing 7 Antibody
Family / Subfamily:	Tripartite Motif
Host	TRIM7 antibody was produced in Goat
Clonality:	Polyclonal
Immunogen Species:	TRIM7 antibody was raised against Human
Antigen Type:	Synthetic peptide
Immunogen:	TRIM7 antibody was raised against synthetic peptide C-QAPAGPPWDITEA from the C-terminus of human TRIM7 (NP_203128.1). Percent identity with other species by BLAST analysis: Human (100%).
Specificity:	Human TRIM7.
Epitope:	C-Terminus
Reactivity:	Human
Purification:	Immunoaffinity purified
Presentation:	Tris-buffered saline, pH 7.3, 0.5% BSA, 0.02% sodium azide
Recommended Storage:	Store at -20°C. Minimize freezing and thawing.
Uses:	IHC - Paraffin (3.75 µg/ml), Western blot (1:32000) & (1 µg/ml), ELISA (1:32000) (Optimal dilution to be determined by the researcher)
Size:	50 µg
Concentration:	0.5 mg/ml

Immunohistochemistry Image:




Anti-TRIM7 antibody IHC of human skeletal muscle. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B4968 concentration 3.75 ug/ml.

Immunohistochemistry Image:




Anti-TRIM7 antibody IHC of human adipocytes. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B4968 concentration 3.75 ug/ml.

Immunohistochemistry Image:



Anti-TRIM7 antibody IHC of human lung. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B4968 concentration 3.75 ug/ml.

Western Blot Image:



Antibody staining (1 ug/ml) of Human Muscle lysate (RIPA buffer, 30 ug total protein per lane). Primary incubated for 1 hour. Detected by Western blot of chemiluminescence.

Requested From:

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/24/2014

© 2014 LifeSpan BioSciences