


**USP10 Rabbit anti-Human Polyclonal (N-Terminus) Antibody - LS-B5278 - LSBio**


<b>CatalogID:</b>	LS-B5278
<b>Validation:</b>	This antibody replaces catalog number LS-C115852. It has been validated for use in the following assays: IHC-P.
<b>Target:</b>	ubiquitin specific peptidase 10 (USP10)
<b>Synonyms:</b>	USP10 Antibody, KIAA0190 Antibody, Ubiquitin thiolesterase 10 Antibody, UBPO Antibody, Ubiquitin specific protease 10 Antibody, Deubiquitinating enzyme 10 Antibody, Ubiquitin thioesterase 10 Antibody
<b>Family / Subfamily:</b>	Protease / Cysteine C19
<b>Host</b>	USP10 antibody was produced in Rabbit
<b>Clonality:</b>	Polyclonal
<b>Immunogen Species:</b>	USP10 antibody was raised against Human
<b>Antigen Type:</b>	Synthetic peptide
<b>Immunogen:</b>	USP10 antibody was raised against a 16 amino acid peptide near the amino terminus of human USP10.
<b>Specificity:</b>	Human USP10
<b>Epitope:</b>	N-Terminus
<b>Reactivity:</b>	Human, Mouse
<b>Purification:</b>	Immunoaffinity purified
<b>Presentation:</b>	PBS, 0.02% sodium azide
<b>Recommended Storage:</b>	Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles. Store undiluted.
<b>Usage Summary:</b>	USP10 antibody can be used for detection of USP10 by Western blot at 1-2 ug/ml.
<b>Uses:</b>	IHC - Paraffin (5 µg/ml), ICC, Western blot (1 - 2 µg/ml), ELISA (Optimal dilution to be determined by the researcher)
<b>Size:</b>	50 µg
<b>Concentration:</b>	1 mg/ml

**Immunohistochemistry Image:**




Anti-USP10 antibody IHC of human breast. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B5278 concentration 5 ug/ml.

**Immunohistochemistry Image:**




Anti-USP10 antibody IHC of human colon. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B5278 concentration 5 ug/ml.

**Immunohistochemistry Image:**



Anti-USP10 antibody IHC of human brain, cortex. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B5278 concentration 5 ug/ml.

**Immunofluorescence Image:**




Immunofluorescence of USP10 in Jurkat cells with USP10 antibody at 20 ug/ml.

**Immunocytochemistry Image:**



Immunocytochemistry of USP10 in Jurkat cells with USP10 antibody at 20 ug/ml.

**Western Blot Image:**



Western blot analysis of USP10 in Jurkat cell lysate with USP10 antibody at (A) 1 and (B) 2 ug/ml.

**Requested From:**

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/24/2014

© 2014 LifeSpan BioSciences