


| CST3 / Cystatin C Goat anti-Human Polyclonal Antibody - LS-B5516 - LSBio |  |
|--|--|
| <b>CatalogID:</b>  | LS-B5516   |
| <b>Validation:</b>   | This antibody replaces catalog number LS-C64164. It has been validated for use in the following assays: IHC-P.   |
| <b>Target:</b>   | cystatin C (CST3)  |
| <b>Synonyms:</b>   | CST3 Antibody, ARMD11 Antibody, BA218C14.4 (cystatin C) Antibody, Cystatin-3 Antibody, Cystatin 3 Antibody, Cystatin C Antibody, Cystatin-C Antibody, Gamma-trace Antibody, Post-gamma-globulin Antibody |
| <b>Host</b>  | CST3 antibody was produced in Goat   |
| <b>Clonality:</b>  | Polyclonal   |
| <b>Immunogen Species:</b>  | CST3 / Cystatin C antibody was raised against Human  |
| <b>Antigen Type:</b>   | Purified protein   |
| <b>Immunogen:</b>  | CST3 / Cystatin C antibody was raised against cystatin C purified from human urine.  |
| <b>Specificity:</b>  | Recognizes human cystatin C.   |
| <b>Reactivity:</b>   | Human  |
| <b>Purification:</b>   | Immunoaffinity purified  |
| <b>Presentation:</b>   | PBS, pH 7.2 0.05% sodium azide.  |
| <b>Recommended Storage:</b>  | May be stored at 4°C for short-term only. Aliquot to avoid repeated freezing and thawing. Store at -20°C. Aliquots are stable for at least 12 months.  |
| <b>Usage Summary:</b>  | Suitable for use in ELISA.   |
| <b>Uses:</b>   | IHC - Paraffin (5 µg/ml), ELISA (Optimal dilution to be determined by the researcher)  |
| <b>Size:</b>   | 50 µg  |
| <b>Concentration:</b>  | 1 mg/ml  |

**Immunohistochemistry Image:**



Anti-CST3 / Cystatin C antibody IHC of human kidney, tubules. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B5516 concentration 5 ug/ml.

**Requested From:**

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/24/2014

© 2014 LifeSpan BioSciences