



BIK Mouse anti-Human Monoclonal (4G12) Antibody - LS-B6011 - LSBio	
CatalogID:	LS-B6011
Validation:	This antibody replaces catalog number LS-C133002. It has been validated for use in the following assays: IHC-P.
Target:	BCL2-interacting killer (apoptosis-inducing) (BIK)
Synonyms:	BIK Antibody, Apoptosis inducer NBK Antibody, Bcl-2-interacting killer Antibody, BP4 Antibody, Apoptosis-inducing NBK Antibody, BIP1 Antibody, NBK Antibody
Host	BIK antibody was produced in Mouse
Clonality:	Monoclonal
Isotype:	IgG
Clone Name:	4G12
Immunogen Species:	BIK antibody was raised against Human
Immunogen:	BIK antibody was raised against bIK (AAH01599, 37 a.a. ~ 136 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
Specificity:	Human Bik
Reactivity:	Human
Purification:	Protein A purified
Presentation:	PBS, pH 7.2
Recommended Storage:	Aliquot and store at -20°C or -80°C. Avoid freeze-thaw cycles.
Usage Summary:	Immunohistochemistry: Formalin-fixed paraffin-embedded sections. Sandwich ELISA: Recombinant protein. Western Blot: Recombinant protein.
Uses:	IHC - Paraffin (5 μg/ml), Western blot (1:500 - 1:1000), ELISA (Optimal dilution to b determined by the researcher)
Size:	50 µg
Concentration:	0.5 mg/ml

Immunohistochemistry Image:



Anti-BIK antibody IHC of human tonsil. Immunohistochemistry of formalin-fixed, paraffinembedded tissue after heat-induced antigen retrieval. Antibody LS-B6011 concentration 5 ug/ml.

ELISA Image:



Recombinant ProteinConcentration(ng/ml)

Detection limit for recombinant GST tagged BIK is approximately 0.03 ng/ml as a capture antibody.

Requested From:	Japan	
Laboratory Reagent For In Vitro Research Use Only		
Not for resale without prior written consent from LifeSpan BioSciences, Inc.		
Created on 9/24/2014		
© 2014 LifeSpan BioSciences		