



TNFRSF12A / FN14 Rabbit anti-Human Polyclonal Antibody - LS-B605 - LSBio	
<b>CatalogID:</b>	LS-B605
<b>Validation:</b>	This antibody replaces catalog number LS-C19643. It has been validated for use in the following assays: IHC.
<b>Target:</b>	tumor necrosis factor receptor superfamily, member 12A (TNFRSF12A)
<b>Synonyms:</b>	TNFRSF12A Antibody, CD266 antigen Antibody, CD266 Antibody, FGF-inducible 14 Antibody, FN14 Antibody, Tweak-receptor Antibody, TWEAKR Antibody
<b>Host</b>	TNFRSF12A antibody was produced in Rabbit
<b>Clonality:</b>	Polyclonal
<b>Immunogen Species:</b>	TNFRSF12A / FN14 antibody was raised against Human
<b>Specificity:</b>	human IgG1 Fc-tagged Fn14 protein
<b>Reactivity:</b>	Human
<b>Purification:</b>	Affinity purified
<b>Presentation:</b>	PBS, 0.02% sodium azide.
<b>Recommended Storage:</b>	Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles. Store undiluted.
<b>Usage Summary:</b>	Immunohistochemistry: LS-B605 was validated for use in immunohistochemistry on a panel of 21 formalin-fixed, paraffin-embedded (FFPE) human tissues after heat induced antigen retrieval in pH 6.0 citrate buffer. After incubation with the primary antibody, slides were incubated with biotinylated secondary antibody, followed by alkaline phosphatase-streptavidin and chromogen. The stained slides were evaluated by a pathologist to confirm staining specificity. The optimal working concentration for LS-B605 was determined to be 2.5 ug/ml.
<b>Uses:</b>	IHC - Paraffin (2) & (2.5 µg/ml), Western blot (1 µg/ml) (Optimal dilution to be determined by the researcher)
<b>Size:</b>	50 µg
<b>Concentration:</b>	1 mg/ml

**Immunohistochemistry Image:**




Anti-TWEAKR antibody IHC of human placenta. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B605 concentration 5 ug/ml.

**Immunohistochemistry Image:**




Anti-TWEAKR antibody IHC of human kidney. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B605 concentration 5 ug/ml.

**Immunofluorescence Image:**



Immunofluorescence of Fn14 in Human Liver cells with Fn14 antibody at 20 ug/ml.

Western Blot Image:



Western blot analysis of Fn14 in HepG2 cells with Fn14 antibody at (A) 2 and (B) 4 ug/ml.

Requested From:

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/24/2014

© 2014 LifeSpan BioSciences