


TLR10 Rabbit Polyclonal Antibody - LS-B625 - LSBio


CatalogID:	LS-B625
Validation:	This antibody replaces catalog number LS-C26901. It has been validated for use in the following assays: IHC.
Target:	toll-like receptor 10 (TLR10)
Synonyms:	TLR10 Antibody, CD290 Antibody, CD290 antigen Antibody, Toll-like receptor 10 Antibody
Family / Subfamily:	Toll-like Receptor / not assigned-Toll-like Receptor
Host	TLR10 antibody was produced in Rabbit
Clonality:	Polyclonal
Antigen Type:	Synthetic peptide
Immunogen:	TLR10 antibody was raised against a peptide corresponding to 14 amino acids near the middle of human TLR10
Reactivity:	Human
Purification:	Purified
Presentation:	PBS, 0.02% sodium azide.
Recommended Storage:	+4°C or -20°C, Avoid repeated freezing and thawing.
Usage Summary:	Immunohistochemistry: LS-B625 was validated for use in immunohistochemistry on a panel of 21 formalin-fixed, paraffin-embedded (FFPE) human tissues after heat induced antigen retrieval in pH 6.0 citrate buffer. After incubation with the primary antibody, slides were incubated with biotinylated secondary antibody, followed by alkaline phosphatase-streptavidin and chromogen. The stained slides were evaluated by a pathologist to confirm staining specificity. The optimal working concentration for LS-B625 was determined to be 2.5 ug/ml.
Uses:	IHC - Paraffin (2.5 µg/ml), Western blot (Optimal dilution to be determined by the researcher)
Size:	50 µg
Concentration:	1 mg/ml

Immunohistochemistry Image:



Anti-TLR10 antibody IHC of human brain, cerebellum. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B625 concentration 5 ug/ml.

Western Blot Image:



Western blot of TLR10 in human lymph node cell lysates with TLR10 antibody (IN) at (A) 0.5 and (B) 1 g/m

Requested From:

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/24/2014

© 2014 LifeSpan BioSciences