



IL8 Mouse anti-Human Monoclonal Antibody - LS-B6427 - LSBio	
CatalogID:	LS-B6427
Validation:	This antibody replaces catalog number LS-C137621. It has been validated for use in the following assays: IHC-P.
Target:	interleukin 8 (IL8)
Synonyms:	IL8 Antibody, 3-10C Antibody, AMCF-I Antibody, B-ENAP Antibody, C-X-C motif chemokine 8 Antibody, GCP-1 Antibody, IL-8 Antibody, Interleukin-8 Antibody, K60 Antibody, LECT Antibody, GCP1 Antibody, LUCT Antibody, NAF Antibody, NAP1 Antibody, Neutrophil-activating factor Antibody, Interleukin 8 Antibody, Protein 3-10C Antibody, MDNCF Antibody, TSG-1 Antibody, CXCL8 Antibody, Emoctakin Antibody, LYNAP Antibody, MONAP Antibody, NAP-1 Antibody, SCYB8 Antibody, T-cell chemotactic factor Antibody
Family / Subfamily:	Intercrine / Intercrine alpha
Host	IL8 antibody was produced in Mouse
Clonality:	Monoclonal
Isotype:	IgG1
Immunogen Species:	IL8 antibody was raised against Human
Antigen Type:	Recombinant protein
Immunogen:	IL8 antibody was raised against recombinant protein
Specificity:	Anti-IL8 recognizes recombinant protein IL8.
Reactivity:	Human
Purification:	Affinity purified
Presentation:	Citrate-Tris-HCl buffer, pH 7.0, 0.02% Proclin 300
Recommended Storage:	Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles.
Uses:	IHC - Paraffin (5 μg/ml), Western blot (1:1000), ELISA (Optimal dilution to be determined by the researcher)
Size:	50 μl
Concentration:	1 mg/ml

## Immunohistochemistry Image:



Anti-IL-8 antibody IHC of human breast, epithelium. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B6427 concentration 5 ug/ml.

## Western Blot Image:



Requested From: Japan