


PPBP / CXCL7 Goat anti-Human Polyclonal Antibody - LS-B7395 - LSBio	
CatalogID:	LS-B7395
Validation:	This antibody replaces catalog number LS-C104406. It has been validated for use in the following assays: IHC-P.
Target:	pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) (PPBP)
Synonyms:	PPBP Antibody, Beta-thromboglobulin Antibody, B-TG1 Antibody, Beta-TG Antibody, C-X-C motif chemokine 7 Antibody, CTAP3 Antibody, CXCL7 Antibody, CTAP-III Antibody, CXC chemokine ligand 7 Antibody, NAP-2 Antibody, NAP-2-L1 Antibody, PBP Antibody, LA-PF4 Antibody, LDGF Antibody, SCYB7 Antibody, Small-inducible cytokine B7 Antibody, Small inducible cytokine B7 Antibody, THBGB1 Antibody, Thrombocidin 2 Antibody, TC2 Antibody, TGB Antibody, Thrombocidin 1 Antibody, Thromboglobulin, beta-1 Antibody, CTAPIII Antibody, MDGF Antibody, Platelet basic protein Antibody, SCAR10 Antibody, TC1 Antibody, THBGB Antibody
Family / Subfamily:	Interocrine / Interocrine alpha
Host	PPBP antibody was produced in Goat
Clonality:	Polyclonal
Isotype:	IgG
Immunogen Species:	PPBP / CXCL7 antibody was raised against Human
Antigen Type:	Cell extract
Immunogen:	PPBP / CXCL7 antibody was raised against e.coli derived recombinant Human NAP-2.
Specificity:	Human NAP-2
Reactivity:	Human
Purification:	Immunoaffinity purified
Reconstitution:	Sterile water. Possible additional volumes for resuspension: 50 µl
Presentation:	Lyophilized from PBS, pH 7.2
Recommended Storage:	Store Lyophilized at room temperature up to 1 month; Reconstituted for up to two weeks at 2-8°C. Aliquot and freeze at -20°C for long term storage. Avoid freeze/thaw cycles.
Usage Summary:	Immunohistochemistry: LS-B7395 in an IHC-plus antibody, validated by our pathologists for use in IHC against formalin-fixed, paraffin-embedded human tissues (IHC validation procedure). A concentration of 5 µg/ml is recommended. ELISA: To detect hNAP-2 by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5-2 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with Biotinylated Anti-Human NAP-2 (LS-C104422) as a detection antibody, allows the detection of at least 0.2-0.4 ng/well of recombinant hNAP-2. Western Blot: To detect hNAP-2 by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hNAP-2 is 1.5-3 ng/lane, under either reducing or non-reducing conditions.
Uses:	IHC - Paraffin (5 µg/ml), Western blot (0.1 - 0.2 µg/ml), ELISA (0.5 - 2 µg/ml) (Optimal dilution to be determined by the researcher)


Size:	50 µg
--------------	-------

Immunohistochemistry Image:




Anti-PPBP antibody IHC of human intravascular platelets. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B7395 concentration 5 ug/ml.

Western Blot Image:




Western Blot (non-reducing) of PPBP / CXCL7 antibody LS-B7395

Western Blot Image:



Western Blot of PPBP / CXCL7 antibody LS-B7395

ELISA Image:



Sandwich ELISA of PPBP / CXCL7 antibody LS-B7395

Requested From:

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/24/2014

© 2014 LifeSpan BioSciences