



FST / Follistatin Rabbit anti-Human Polyclonal Antibody - LS-B7434 - LSBio	
CatalogID:	LS-B7434
Validation:	This antibody replaces catalog number LS-C104466. It has been validated for use in the following assays: IHC-P.
Target:	follistatin (FST)
Synonyms:	FST Antibody, Activin-binding protein Antibody, Follistatin isoform FST317 Antibody, Follistatin Antibody, Fs Antibody
Host	FST antibody was produced in Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Immunogen Species:	FST / Follistatin antibody was raised against Human
Antigen Type:	Cell extract
Immunogen:	FST / Follistatin antibody was raised against e.coli derived recombinant Human Follistatin.
Specificity:	Human Follistatin
Reactivity:	Human
Purification:	Immunoaffinity purified
Reconstitution:	Sterile water. Possible additional volumes for resuspension: 50 µl
Presentation:	Lyophilized from PBS, pH 7.2
Recommended Storage:	Store Lyophilized at room temperature up to 1 month; Reconstituted for up to two weeks at 2-8°C. Aliquot and freeze at -20°C for long term storage. Avoid freeze/thaw cycles.
Usage Summary:	Immunohistochemistry: LS-B7434 in an IHC-plus antibody, validated by our pathologists for use in IHC against formalin-fixed, paraffin-embedded human tissues (IHC validation procedure). A concentration of 5 µg/ml is recommended. Neutralization: To yield one-half maximal inhibition [ND50] of the biological activity of Human Follistatin (100 ng/ml), a concentration of 8-12 µg/ml of this antibody is required. ELISA: To detect Human Follistatin by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5-2 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with Biotinylated Anti-Human Follistatin (LS-C104465) as a detection antibody, allows the detection of at least 0.2-0.4 ng/well of recombinant Human Follistatin. Western Blot: To detect Human Follistatin by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. When used in conjunction with compatible secondary reagents the detection limit for recombinant Human Follistatin is 1.5-3 ng/lane, under either reducing or non-reducing conditions.
Uses:	IHC - Paraffin (5 µg/ml), Western blot (0.1 - 0.2 µg/ml), ELISA (0.5 - 2 µg/ml), Neutralization (8 - 12 µg/ml) (Optimal dilution to be determined by the researcher)
Size:	50 µg

Immunohistochemistry Image:




Anti-FST / Follistatin antibody IHC of human heart. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B7434 concentration 5 ug/ml.

Western Blot Image:




Western Blot (reducing) of Follistatin antibody LS-B7434

Western Blot Image:



Western Blot (non-reducing) of Follistatin antibody LS-B7434

ELISA Image:



Sandwich ELISA of Follistatin antibody LS-B7434

Requested From:

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/24/2014

© 2014 LifeSpan BioSciences