


IL22 Rabbit anti-Human Polyclonal Antibody - LS-B7437 - LSBio


CatalogID:	LS-B7437
Validation:	This antibody replaces catalog number LS-C104291. It has been validated for use in the following assays: IHC-P.
Target:	interleukin 22 (IL22)
Synonyms:	IL22 Antibody, Cytokine Zcyto18 Antibody, ILTIF Antibody, Interleukin 22 Antibody, IL-22 Antibody, IL-D110 Antibody, TIFIL-23 Antibody, Zcyto18 Antibody, IL-TIF Antibody, Interleukin-22 Antibody
Family / Subfamily:	Interleukin
Host	IL22 antibody was produced in Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Immunogen Species:	IL22 antibody was raised against Human
Antigen Type:	Cell extract
Immunogen:	IL22 antibody was raised against e.coli derived recombinant Human IL-22.
Specificity:	Human IL-22
Reactivity:	Human
Purification:	Immunoaffinity purified
Reconstitution:	Sterile water. Possible additional volumes for resuspension: 50 µl
Presentation:	Lyophilized from PBS, pH 7.2
Recommended Storage:	Store Lyophilized at room temperature up to 1 month; Reconstituted for up to two weeks at 2-8°C. Aliquot and freeze at -20°C for long term storage. Avoid freeze/thaw cycles.
Usage Summary:	Immunohistochemistry: LS-B7437 in an IHC-plus antibody, validated by our pathologists for use in IHC against formalin-fixed, paraffin-embedded human tissues (IHC validation procedure). A concentration of 5 ug/ml is recommended. Neutralization: To yield one-half maximal inhibition [ND] of the biological activity of Human IL-22 (50 ng/ml), a concentration of 1.25 ug/ml of this antibody is required. ELISA: To detect hIL-22 by sandwich ELISA (using 100 ul/well antibody solution) a concentration of 0.5-2 ug/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with Biotinylated Anti-Human IL-22 (LS-C104319) as a detection antibody, allows the detection of at least 0.2-0.4 ng/well of recombinant hIL-22. Western Blot: To detect hIL-22 by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 ug/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hIL-22 is 1.5-3 ng/lane, under either reducing or non-reducing conditions.
Uses:	IHC - Paraffin (5 µg/ml), Western blot (0.1 - 0.2 µg/ml), ELISA (0.5 - 2 µg/ml), Neutralization (1.25 µg/ml) (Optimal dilution to be determined by the researcher)
Size:	50 µg

Immunohistochemistry Image:



Anti-IL-22 antibody IHC of human tonsil. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B7437 concentration 5 ug/ml.

ELISA Image:



Sandwich ELISA of IL-22 antibody LS-B7437

Requested From:

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/24/2014

© 2014 LifeSpan BioSciences