


| TNF Alpha Rabbit anti-Human Polyclonal Antibody - LS-B7730 - LSBio |   |
|--|---|
| <b>CatalogID:</b>  | LS-B7730  |
| <b>Validation:</b>   | This antibody replaces catalog number LS-C48163. It has been validated for use in the following assays: IHC-P.  |
| <b>Target:</b>   | tumor necrosis factor (TNF)   |
| <b>Synonyms:</b>   | TNF Antibody, APC1 protein Antibody, Cachectin Antibody, DIF Antibody, TNF Alpha Antibody, Tumor necrosis factor-alpha Antibody, TNF-alpha Antibody, TNF-a Antibody, TNFA Antibody, Tumor necrosis factor Antibody, TNF, macrophage-derived Antibody, TNF, monocyte-derived Antibody, TNFSF2 Antibody |
| <b>Family / Subfamily:</b>   | TNF / not assigned-TNF  |
| <b>Host</b>  | TNF antibody was produced in Rabbit   |
| <b>Clonality:</b>  | Polyclonal  |
| <b>Immunogen Species:</b>  | TNF Alpha antibody was raised against Human   |
| <b>Antigen Type:</b>   | Recombinant protein   |
| <b>Immunogen:</b>  | TNF Alpha antibody was raised against e. coli expressed recombinant human TNF -alpha.   |
| <b>Reactivity:</b>   | Human   |
| <b>Purification:</b>   | Affinity purified   |
| <b>Presentation:</b>   | PBS, pH 7.2, 30% glycerol, 0.5% BSA, 0.01% Thimerosal.  |
| <b>Recommended Storage:</b>  | Long term: -70°C; Short term: -20°C   |
| <b>Uses:</b>   | IHC - Paraffin (5 µg/ml), Western blot (0.5 - 4 µg/ml), Neutralization (0.5 - 5 µg/ml) (Optimal dilution to be determined by the researcher)  |
| <b>Size:</b>   | 50 µg   |
| <b>Concentration:</b>  | 0.5 mg/ml   |

**Immunohistochemistry Image:**



Anti-TNF Alpha / TNF antibody IHC of human tonsil. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B7730 dilution 5 ug/ml.

**Requested From:**

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/24/2014

© 2014 LifeSpan BioSciences