


RCOR1 / COREST Rabbit anti-Human Polyclonal (aa109-293) Antibody - LS-B8140 - LSBio	
<b>CatalogID:</b>	LS-B8140
<b>Validation:</b>	This antibody replaces catalog number LS-C14592. It has been validated for use in the following assays: IHC-P.
<b>Target:</b>	REST corepressor 1 (RCOR1)
<b>Synonyms:</b>	RCOR1 Antibody, COREST Antibody, Protein CoREST Antibody, REST corepressor 1 Antibody, RCOR Antibody, KIAA0071 Antibody, REST corepressor Antibody
<b>Host</b>	RCOR1 antibody was produced in Rabbit
<b>Clonality:</b>	Polyclonal
<b>Isotype:</b>	IgG
<b>Immunogen Species:</b>	RCOR1 / COREST antibody was raised against Human
<b>Antigen Type:</b>	Fusion protein
<b>Immunogen:</b>	RCOR1 / COREST antibody was raised against gST fusion protein corresponding to residues 109-293 of human CoREST.
<b>Specificity:</b>	Recognizes human CoREST, Mr ~66kD. Species cross-reactivity: Mouse and rat.
<b>Epitope:</b>	aa109-293
<b>Reactivity:</b>	Human, Mouse, Rat
<b>Purification:</b>	Protein A purified
<b>Presentation:</b>	0.1 M Tris-glycine, pH 7.4, 0.15 M sodium chloride, 0.05% sodium azide, 30% glycerol.
<b>Recommended Storage:</b>	Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles. Store undiluted.
<b>Usage Summary:</b>	Suitable for use in Western Blot, Immunoprecipitation, Immunohistochemistry, Immunocytochemistry. Add 2.5 ul of 2-mercaptoethanol/100 ul of lysate and boil for 5 minutes to reduce the preparation. Load 20 ug of reduced lysate per lane for minigels. Positive control: A431 Cells, Non-Stimulated in RIPA Buffer, Sample Buffer. Add 2.5 ul of 2-mercaptoethanol/100 ul of lysate and boil for 5 minutes to reduce the preparation. Load 20 ug of reduced lysate per lane for minigels.
<b>Uses:</b>	IHC - Paraffin (10 µg/ml), ICC, Western blot, Immunoprecipitation (Optimal dilution to be determined by the researcher)
<b>Size:</b>	50 µg
<b>Concentration:</b>	1 mg/ml

**Immunohistochemistry Image:**



Anti-COREST antibody IHC of human breast. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B8140 dilution 10 ug/ml.

**Requested From:**

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/24/2014

© 2014 LifeSpan BioSciences