


STAD2 Cost anti Human Dalvalanal (C Terminus) Antibody, LS D2202, LSDie	
STAP2 Goat anti-Human Polyclonal (C-Terminus) Antibody - LS-B8288 - LSBio	
CatalogID:	LS-B8288
Validation:	This antibody replaces catalog number LS-C112902. It has been validated for use in the following assays: IHC-P.
Target:	signal transducing adaptor family member 2 (STAP2)
Synonyms:	STAP2 Antibody, BRK substrate Antibody, BKS Antibody, Brk kinase substrate Antibody, STAP-2 Antibody, Breast tumor kinase substrate Antibody
Host	STAP2 antibody was produced in Goat
Clonality:	Polyclonal
Immunogen Species:	STAP2 antibody was raised against Human
Antigen Type:	Synthetic peptide
Immunogen:	STAP2 antibody was raised against synthetic peptide C-LQKKLEKRRALEH from the C-terminus of human STAP2 (NP_060190.2; NP_001013863.1). Percent identity by BLAST analysis: Human, Gorilla, Gibbon, Marmoset (100%).
Specificity:	Human STAP2. This antibody is expected to recognize isoform 1 (NP_060190.2) and isoform 2 (NP_001013863.1).
Epitope:	C-Terminus
Reactivity:	Human, Gorilla, Gibbon, Monkey
Purification:	Immunoaffinity purified
Presentation:	Tris-buffered saline, pH 7.3, 0.5% BSA, 0.02% sodium azide
Recommended Storage:	Store at -20°C. Minimize freezing and thawing.
Usage Summary:	Peptide ELISA: antibody detection limit dilution 1:128000. Western blot: Approx 49kDa band observed in Human Breast lysates (calculated MW of 49.6kDa according to NP_060190.2). Recommended concentration: 0.3-1 ug/ml.
Uses:	IHC - Paraffin (3.75 μg/ml), Western blot (0.3 - 1 μg/ml), ELISA (1:128000) (Optimal dilution to be determined by the researcher)
Size:	50 μg
Concentration:	0.5 mg/ml

Immunohistochemistry Image:



Anti-STAP2 antibody IHC of human small intestine. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B8288 dilution 3.75 ug/ml.

Western Blot Image:

250kDa 150kDa 100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

15kDa

STAP2 antibody (0.3 ug/ml) staining of Human Breast lysate (35 ug protein/ml in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Requested From: Japan

Laboratory Reagent For In Vitro Research Use Only
Not for resale without prior written consent from LifeSpan BioSciences, Inc.
Created on 9/24/2014
© 2014 LifeSpan BioSciences