


**ST13 Rabbit anti-Human Polyclonal (aa60-75) Antibody - LS-B903 - LSBio**

<b>CatalogID:</b>	LS-B903
<b>Validation:</b>	This antibody replaces catalog number LS-C3356. It has been validated for use in the following assays: IHC.
<b>Target:</b>	suppression of tumorigenicity 13 (colon carcinoma) (Hsp70 interacting protein) (ST13)
<b>Synonyms:</b>	ST13 Antibody, Aging-associated protein 2 Antibody, AAG2 Antibody, Hsc70-interacting protein Antibody, HSPABP1 Antibody, HOP Antibody, Hsp70-interacting protein Antibody, HSPABP Antibody, FAM10A1 Antibody, HIP Antibody, p48 Antibody, PRO0786 Antibody, Protein FAM10A1 Antibody, Putative tumor suppressor ST13 Antibody, SNC6 Antibody, FAM10A4 Antibody
<b>Host</b>	ST13 antibody was produced in Rabbit
<b>Clonality:</b>	Polyclonal
<b>Immunogen Species:</b>	ST13 antibody was raised against Human
<b>Antigen Type:</b>	Synthetic peptide
<b>Immunogen:</b>	ST13 antibody was raised against synthetic peptide from human ST13 / HIP.
<b>Specificity:</b>	Amino acids 60 to 75 of human ST13
<b>Epitope:</b>	aa60-75
<b>Reactivity:</b>	Human
<b>Purification:</b>	Protein G purified
<b>Presentation:</b>	PBS, 0.1% sodium azide.
<b>Recommended Storage:</b>	Long term: -20°C; Short term: +4°C. Avoid repeat freeze-thaw cycles.
<b>Usage Summary:</b>	Immunohistochemistry: LS-B903 was validated for use in immunohistochemistry on a panel of 21 formalin-fixed, paraffin-embedded (FFPE) human tissues after heat induced antigen retrieval in pH 6.0 citrate buffer. After incubation with the primary antibody, slides were incubated with biotinylated secondary antibody, followed by alkaline phosphatase-streptavidin and chromogen. The stained slides were evaluated by a pathologist to confirm staining specificity. The optimal working concentration for LS-B903 was determined to be 10 ug/ml.
<b>Uses:</b>	IHC - Paraffin (10 µg/ml), ELISA (1:000 - 1:1000) (Optimal dilution to be determined by the researcher)
<b>Size:</b>	50 µg
<b>Concentration:</b>	1 mg/ml

**Immunohistochemistry Image:**



Anti-ST13 / HIP antibody IHC of human adrenal. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody LS-B903 concentration 10 ug/ml.

**Requested From:**

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/24/2014

© 2014 LifeSpan BioSciences