



MAC-2-BP / LGALS3BP Mouse anti-Human Monoclonal (2B1) Antibody - LS-B9815 - LSBio	
<b>CatalogID:</b>	LS-B9815
<b>Validation:</b>	This antibody replaces catalog number LS-C173265. It has been validated for use in the following assays: IHC-P.
<b>Target:</b>	lectin, galactoside-binding, soluble, 3 binding protein (LGALS3BP)
<b>Synonyms:</b>	LGALS3BP Antibody, 90K Antibody, Galectin-3-binding protein Antibody, Mac-2 BP Antibody, Mac-2-binding protein Antibody, MAC2BP Antibody, L3 antigen Antibody, Serum protein 90K Antibody, TANGO10B Antibody, BTBD17B Antibody, M2BP Antibody, MAC-2-BP Antibody, Tumor-associated antigen 90K Antibody
<b>Host</b>	LGALS3BP antibody was produced in Mouse
<b>Clonality:</b>	Monoclonal
<b>Isotype:</b>	IgG1
<b>Clone Name:</b>	2B1
<b>Immunogen Species:</b>	MAC-2-BP / LGALS3BP antibody was raised against Human
<b>Antigen Type:</b>	Recombinant protein
<b>Immunogen:</b>	MAC-2-BP / LGALS3BP antibody was raised against human recombinant protein fragment corresponding to amino acids 19-300 of human LGALS3BP(NP_005558) produced in E. coli.
<b>Specificity:</b>	Human MAC-2-BP / LGALS3BP
<b>Reactivity:</b>	Human
<b>Purification:</b>	Protein A/G purified
<b>Presentation:</b>	PBS, pH 7.3, 1% BSA, 50% glycerol, 0.02% sodium azide
<b>Recommended Storage:</b>	Store at -20°C. Minimize freezing and thawing.
<b>Uses:</b>	IHC - Paraffin (10 µg/ml), Western blot (1:2000), Flow Cytometry (1:100) (Optimal dilution to be determined by the researcher)
<b>Size:</b>	50 µl
<b>Concentration:</b>	0.81 mg/ml

**Immunohistochemistry Image:**




Human Small Intestine: Formalin-Fixed, Paraffin-Embedded (FFPE)

**Flow Cytometry Image:**



HEK293T cells transfected with either overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-LGALS3BP antibody, and then analyzed by flow cytometry.

**Western Blot Image:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY LGALS3BP (Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-LGALS3BP.

**Requested From:**

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/24/2014

© 2014 LifeSpan BioSciences