


**CD1C Mouse anti-Human Monoclonal (FITC) (L161) Antibody - LS-C106298 - LSBio**

<b>CatalogID:</b>	LS-C106298
<b>Target:</b>	CD1c molecule
<b>Synonyms:</b>	CD1C Antibody, BDCA1 Antibody, CD1 Antibody, CD1c molecule Antibody, CD1C antigen, c polypeptide Antibody, CD1c antigen Antibody, R7 Antibody
<b>Host</b>	CD1C antibody was produced in Mouse
<b>Clonality:</b>	Monoclonal
<b>Isotype:</b>	IgG1
<b>Clone Name:</b>	L161
<b>Conjugations:</b>	Fluorescein (FITC)
<b>Immunogen Species:</b>	CD1C antibody was raised against Human
<b>Immunogen:</b>	CD1C antibody was raised against human CD1c
<b>Reactivity:</b>	Human
<b>Purification:</b>	Affinity purified
<b>Presentation:</b>	PBS, pH 7.2, <=0.09% sodium azide
<b>Recommended Storage:</b>	Store at +4°C. Do not freeze. Product is photosensitive and should be protected from light.
<b>Usage Summary:</b>	This L161 antibody has been pre-titrated and tested by flow cytometric analysis on normal human peripheral blood cells. This can be used at 5 ul (0.06 ug) per test. A test is defined as the amount (ug) of antibody that will stain a cell sample in a final volume of 100 ul. Cell number should be determined empirically but can range from 10 <sup>5</sup> to 10 <sup>8</sup> cells/test.
<b>Uses:</b>	Immunoprecipitation, Flow Cytometry (Optimal dilution to be determined by the researcher)
<b>Size:</b>	25 tst or 100 tst

**Flow Cytometry Image:**



Staining of normal human peripheral blood cells with FITC Mouse IgG1 Isotype Control (left) or anti-human CD1c (right) and APC anti-human CD19 (HIB19). Total viable cells were used for this analysis.

**Requested From:**

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/24/2014

© 2014 LifeSpan BioSciences