


CD27 Hamster anti-Mouse Monoclonal (FITC) (LG.7F9) Antibody - LS-C106329 - LSBio	
CatalogID:	LS-C106329
Target:	CD27 molecule
Synonyms:	CD27 Antibody, CD27 antigen Antibody, CD27 molecule Antibody, S152 Antibody, T14 Antibody, T-cell activation antigen CD27 Antibody, CD27L receptor Antibody, T cell activation antigen S152 Antibody, TNFRSF7 Antibody, Tp55 Antibody
Family / Subfamily:	TNF Receptor / not assigned-TNF Receptor
Host	CD27 antibody was produced in Hamster
Clonality:	Monoclonal
Isotype:	IgG
Clone Name:	LG.7F9
Conjugations:	Fluorescein (FITC)
Immunogen Species:	CD27 antibody was raised against Mouse
Immunogen:	CD27 antibody was raised against mouse CD27
Reactivity:	Mouse
Purification:	Affinity purified
Presentation:	PBS, pH 7.2, 150 mM sodium chloride, 0.09% sodium azide
Recommended Storage:	Store at +4°C. Do not freeze. Product is photosensitive and should be protected from light.
Usage Summary:	The LG.7F9 antibody has been tested by flow cytometric analysis of mouse splenocyte cell suspensions. This can be used at less than or equal to 0.5 ug per test. A test is defined as the amount (ug) of antibody that will stain a cell sample in a final volume of 100 ul. Cell number should be determined empirically but can range from 10 ⁵ to 10 ⁸ cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.
Uses:	Flow Cytometry (Optimal dilution to be determined by the researcher)
Size:	50 µg or 100 µg or 500 µg

Flow Cytometry Image:



Staining of C57Bl/6 splenocytes with APC anti-mo/hu B220 (RA3-6B2) (LS-C107315) and 0.25 ug of FITC Armenian Hamster IgG isotype control (left) or 0.25 ug of FITC anti-mo/hu/rat CD27 (LG.7F9) (right). Total viable cells were used for analysis.

Requested From:

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/24/2014

© 2014 LifeSpan BioSciences