


**CD24 Rat anti-Mouse Monoclonal (FITC) (M1/69) Antibody - LS-C106332 - LSBio**

<b>CatalogID:</b>	LS-C106332
<b>Target:</b>	CD24 molecule
<b>Synonyms:</b>	CD24 Antibody, CD24 molecule Antibody, CD24A Antibody, Signal transducer CD24 Antibody, CD24 antigen Antibody
<b>Host</b>	CD24 antibody was produced in Rat
<b>Clonality:</b>	Monoclonal
<b>Isotype:</b>	IgG2b,k
<b>Clone Name:</b>	M1/69
<b>Conjugations:</b>	Fluorescein (FITC)
<b>Immunogen Species:</b>	CD24 antibody was raised against Mouse
<b>Immunogen:</b>	CD24 antibody was raised against mouse CD24
<b>Reactivity:</b>	Mouse
<b>Purification:</b>	Affinity purified
<b>Presentation:</b>	PBS, pH 7.2, 150 mM sodium chloride, 0.09% sodium azide
<b>Recommended Storage:</b>	Store at +4°C. Do not freeze. Product is photosensitive and should be protected from light.
<b>Usage Summary:</b>	The M1/69 antibody has been tested by flow cytometric analysis of mouse splenocytes and thymocytes. This can be used at less than or equal to 0.5 ug per test. A test is defined as the amount (ug) of antibody that will stain a cell sample in a final volume of 100 ul. Cell number should be determined empirically but can range from 10 <sup>5</sup> to 10 <sup>8</sup> cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.
<b>Uses:</b>	Flow Cytometry (Optimal dilution to be determined by the researcher)
<b>Size:</b>	50 µg or 100 µg or 500 µg

**Flow Cytometry Image:**



Staining of C57Bl/6 splenocytes with APC anti-mouse CD3e (145-2C11) (LS-C107273) and 0.25 ug of FITC Rat IgG2b Isotype Control (left) or 0.25 ug of FITC anti-mouse CD24 (M1/69) (right). Cells in the lymphocyte gate were used for analysis.

**Requested From:**

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/24/2014

© 2014 LifeSpan BioSciences