



FCER2 / CD23 Mouse anti-Human Monoclonal (FITC) (EBVCS2) Antibody - LS-C106334 - LSBio	
CatalogID:	LS-C106334
Target:	Fc fragment of IgE, low affinity II, receptor for (CD23) (FCER2)
Synonyms:	FCER2 Antibody, CD23 antigen Antibody, CD23A Antibody, CLEC4J Antibody, CD23 Antibody, Fc-epsilon-RII Antibody, IGEBF Antibody, Lymphocyte IgE receptor Antibody, BLAST-2 Antibody, FCE2 Antibody
Host	FCER2 antibody was produced in Mouse
Clonality:	Monoclonal
Isotype:	IgG1
Clone Name:	EBVCS2
Conjugations:	Fluorescein (FITC)
Immunogen Species:	FCER2 / CD23 antibody was raised against Human
Immunogen:	FCER2 / CD23 antibody was raised against human FCER2
Reactivity:	Human
Purification:	Affinity purified
Presentation:	PBS, pH 7.2, 150 mM sodium chloride, 0.09% sodium azide, 0.2% BSA
Recommended Storage:	Store at +4°C. Do not freeze. Product is photosensitive and should be protected from light.
Usage Summary:	The EBVCS2 antibody has been pre-titrated and tested by flow cytometric analysis of human peripheral blood leukocytes. This can be used at 5 ul (0.25 ug)/per test. A test is defined as the amount (ug)/test of antibody that will stain a cell sample in a final volume of 100 ul. Cell number should be determined empirically but can range from 10 ⁵ to 10 ⁸ cells/test.
Uses:	IHC - Frozen, Flow Cytometry (Optimal dilution to be determined by the researcher)
Size:	25 tst or 100 tst

Flow Cytometry Image:



Staining of normal human peripheral blood cells with APC anti-human CD19 (HIB19) and 0.125 ug of FITC Mouse IgG1, K isotype control. Cells in the lymphocyte gate were used for analysis.

Flow Cytometry Image:



Staining of normal human peripheral blood cells with APC anti-human CD19 (HIB19) and 0.25 ug of FITC EBVCS2. Cells in the lymphocyte gate were used for analysis.

Requested From:

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/24/2014

© 2014 LifeSpan BioSciences