

CatalogID:	LS-C112540
Target:	CD79a molecule, immunoglobulin-associated alpha
Synonyms:	CD79A Antibody, CD79a antigen Antibody, Ig-alpha Antibody, MB-1 Antibody, MB- 1 membrane glycoprotein Antibody, Surface IgM-associated protein Antibody, MB1 Antibody
Host	CD79A antibody was produced in Mouse
Clonality:	Monoclonal
Isotype:	IgG1
Clone Name:	HM57
Conjugations:	Phycoerythrin (PE)
Immunogen Species:	CD79A antibody was raised against Human
Antigen Type:	Synthetic peptide - KLH conjugated
Immunogen:	CD79A antibody was raised against synthetic peptide corresponding to amino acids 202-216 of human CD79a.
Specificity:	The antibody HM57 interacts with CD79a (Ig alpha), a 40-45 kD subunit of B cell antigen-specific receptor (BCR) and its early developmental forms. HLDA V; WS Code BC cB018. HLDA VI; WS Code BP 193. HLDA VI; WS Code BP 89. HLDA VI; WS Code B B103. HLDA VI; WS Code B CD79.4.
Reactivity:	Human, Mouse, Rat, Bovine, Guinea pig, Horse, Opossum, Pig, Rabbit, Chicken
Purification:	Protein A purified
Presentation:	15 mM sodium azide, 0.2% BSA, PBS
Recommended Storage:	Store at +4°C. Do not freeze. Product is photosensitive and should be protected from light.
Usage Summary:	The reagent is designed for Flow Cytometry analysis of human blood cells using 20 ul reagent / 100 ul of whole blood or 10^6 cells in a suspension. The content of a vial (2 ml) is sufficient for 100 tests.
Uses:	Flow Cytometry (Optimal dilution to be determined by the researcher)
Size:	100 tst
Requested From:	Japan
La	boratory Reagent For In Vitro Research Use Only
Not for resale	without prior written consent from LifeSpan BioSciences, Inc.
	Created on 9/24/2014