

CatalogID:	LS-C112541
Target:	CD86 molecule
Synonyms:	CD86 Antibody, Activation B7-2 antigen Antibody, B7-2 Antibody, B7.2 Antibody, B70 Antibody, CD86 molecule Antibody, CTLA-4 counter-receptor B7.2 Antibody, CD86 antigen Antibody, FUN-1 Antibody, LAB72 Antibody, BU63 Antibody, CD28LG2 Antibody, Ctla-4 counter-receptors b7.2 Antibody
Host	CD86 antibody was produced in Mouse
Clonality:	Monoclonal
Isotype:	IgG1
Clone Name:	BU63
Conjugations:	Phycoerythrin (PE)
Immunogen Species:	CD86 antibody was raised against Human
Immunogen:	CD86 antibody was raised against b-lymphoblastoid cell line ARH 77.
Specificity:	The antibody BU63 reacts with CD86 (B7-2), a 70 kD type I transmembrane glycoprotein of immunoglobulin supergene family, expressed on professional antigen-presenting cells, such as dendritic cells, macrophages or activated B lymphocytes. HLDA V; WS Code BP BP072. HLDA V; WS Code A A109. HLDA VI; WS Code BP 95. HLDA VI; WS Code B CD86.9.
Reactivity:	Human
Purification:	Protein A purified
Presentation:	15 mM sodium azide, 0.2% BSA, PBS
Recommended Storage:	Store at +4°C. Do not freeze. Product is photosensitive and should be protected from light.
Usage Summary:	The reagent is designed for Flow Cytometry analysis of human blood cells using 20 ul reagent / 100 ul of whole blood or 10^6 cells in a suspension. The content of a vial (2 ml) is sufficient for 100 tests.
Uses:	Flow Cytometry (Optimal dilution to be determined by the researcher)
Size:	100 tst
Requested From:	Japan
La	boratory Reagent For In Vitro Research Use Only
Not for resale	without prior written consent from LifeSpan BioSciences, Inc.
	Created on 9/24/2014