

LS-C112549 CD28 molecule CD28 Antibody, CD28 antigen Antibody, CD28 antigen (Tp44) Antibody, Tp44 Antibody, CD28 molecule Antibody CD28 antibody was produced in Mouse Monoclonal IgG1 CD28.2 Phycoerythrin (PE)
CD28 Antibody, CD28 antigen Antibody, CD28 antigen (Tp44) Antibody, Tp44 Antibody, CD28 molecule Antibody CD28 antibody was produced in Mouse Monoclonal IgG1 CD28.2
Antibody, CD28 molecule Antibody CD28 antibody was produced in Mouse Monoclonal IgG1 CD28.2
Monoclonal IgG1 CD28.2
lgG1 CD28.2
CD28.2
Phycoerythrin (PE)
CD28 antibody was raised against Human
CD28 antibody was raised against dC28.1.3.3 hybridoma cells.
The antibody CD28.2 reacts with CD28, a disulfide-linked homodimeric type I glycoprotein (monomer of Mw 44 kD) which is a critical costimulatory receptor of T cells. HLDA V.; WS Code 5T CD28.05.
Human, Primate
Protein A purified
15 mM sodium azide, 0.2% BSA, PBS
Store at +4°C. Do not freeze. Product is photosensitive and should be protected from light.
The reagent is designed for Flow Cytometry analysis of human blood cells using 20 ul reagent / 100 ul of whole blood or 10^6 cells in a suspension. The content of a vial (2 ml) is sufficient for 100 tests.
Flow Cytometry (Optimal dilution to be determined by the researcher)
100 tst
Japan
atory Reagent For In Vitro Research Use Only
hout prior written consent from LifeSpan BioSciences, Inc.
Created on 9/24/2014