

herin 1, type 1, E-cadherin (epithelial) (CDH1) 11 Antibody, CD324 Antibody, CDHE Antibody, Cell-CAM 120/80 Antibody, E herin Antibody, E-Cadherin Antibody, ECAD Antibody, Epithelial cadherin body, Arc-1 Antibody, UVO Antibody, Cadherin-1 Antibody, CAM 120/80 body, CD324 antigen Antibody, LCAM Antibody, Uvomorulin Antibody herin / not assigned-Cadherin 11 antibody was produced in Mouse oclonal 11 / E Cadherin antibody was raised against Human ein 11 / E Cadherin antibody was raised against cadherin E (E-Cadherin, L-CAM, helial Cell Adhesion Molecule) T-47D breast carcinoma line.
herin Antibody, E-Cadherin Antibody, ECAD Antibody, Epithelial cadherin body, Arc-1 Antibody, UVO Antibody, Cadherin-1 Antibody, CAM 120/80 body, CD324 antigen Antibody, LCAM Antibody, Uvomorulin Antibody herin / not assigned-Cadherin I1 antibody was produced in Mouse oclonal I1 / E Cadherin antibody was raised against Human ein I1 / E Cadherin antibody was raised against cadherin E (E-Cadherin, L-CAM,
I1 antibody was produced in Mouse oclonal I1 / E Cadherin antibody was raised against Human ein I1 / E Cadherin antibody was raised against cadherin E (E-Cadherin, L-CAM,
oclonal I1 / E Cadherin antibody was raised against Human ein I1 / E Cadherin antibody was raised against cadherin E (E-Cadherin, L-CAM,
I1 / E Cadherin antibody was raised against Human ein I1 / E Cadherin antibody was raised against cadherin E (E-Cadherin, L-CAM,
 I1 / E Cadherin antibody was raised against Human ein I1 / E Cadherin antibody was raised against cadherin E (E-Cadherin, L-CAM,
ein I1 / E Cadherin antibody was raised against cadherin E (E-Cadherin, L-CAM,
11 / E Cadherin antibody was raised against cadherin E (E-Cadherin, L-CAM,
11 / E Cadherin antibody was raised against cadherin E (E-Cadherin, L-CAM, nelial Cell Adhesion Molecule) T-47D breast carcinoma line.
ognizes human Cadherin E, a 123kD molecule.
an
ein A purified
, pH 7.6, 0.1% sodium azide.
g term: -20°C; Short term: +4°C. Avoid repeat freeze-thaw cycles.
tern Blot FC Analysis Functional blocking of E-cadherin Does not work well for unoprecipitation. Suitable to study E-cadherin expression by flow cytometry.
tern blot, Immunoprecipitation, Flow Cytometry, Functional Assay (Optimal on to be determined by the researcher)
hð
in
Reagent For In Vitro Research Use Only