

CatalogID:	LS-C127295
Target:	chemokine (C-C motif) receptor 5 (gene/pseudogene) (CCR5)
Synonyms:	CCR5 Antibody, C-C chemokine receptor 5 Antibody, CD195 antigen Antibody, CD195 Antibody, C-C CKR-5 Antibody, Chemokine c-c motif receptor 5 Antibody, Chemr13 Antibody, CMKBR5 Antibody, CKR5 Antibody, Cc chemokine receptor 5 Antibody, CCCKR5 Antibody, CCR-5 Antibody, Ccr5 receptor Antibody, Chemokine receptor CCR5 Antibody, HIV-1 fusion coreceptor Antibody, IDDM22 Antibody, C-C chemokine receptor type 5 Antibody, CC-CKR-5 Antibody, CKR-5 Antibody
Family / Subfamily:	GPCR / Chemokine
Host	CCR5 antibody was produced in Mouse
Clonality:	Monoclonal
sotype:	IgG2a
Clone Name:	HEK/1/85A
Conjugations:	Phycoerythrin (PE)
mmunogen Species:	CCR5 antibody was raised against Human
Antigen Type:	Cells
mmunogen:	CCR5 antibody was raised against cHO cells transfected with human CCR5.
Specificity:	Recognizes human CD195 at ~45kD.
Reactivity:	Human
Purification:	Purified
Presentation:	PBS, 0.9% sodium azide, 1% BSA
Recommended Storage:	May be stored at 4°C before opening. DO NOT FREEZE! Stable at 4°C as an undiluted liquid. Dilute only prior to immediate use. Stable for at least 12 months at 4°C. Sensitive to light.
Usage Summary:	Suitable for use in Flow cytometry: 10 ul labels of a neat-1:5 working dilution to 10^6 cells in 100ul.
Uses:	Flow Cytometry (Optimal dilution to be determined by the researcher)
Size:	100 tst
Requested From:	Japan
La	boratory Reagent For In Vitro Research Use Only
Not for resale	without prior written consent from LifeSpan BioSciences, Inc.