

| | anti-Bovine Monoclonal (FITC) Antibody - LS-C127685 - LSBio |
|--|---|
| CatalogID: | LS-C127685 |
| Target: | CD44 molecule (Indian blood group) |
| Synonyms: | CD44 Antibody, CDW44 Antibody, CD44 antigen Antibody, Cell surface glycoprotein CD44 Antibody, CSPG8 Antibody, ECMR-III Antibody, Epican Antibody, HUTCH-I Antibody, LHR Antibody, MDU2 Antibody, MC56 Antibody, MIC4 Antibody, Hermes antigen Antibody, Hyaluronate receptor Antibody, Phagocytic glycoprotein 1 Antibody, PGP-1 Antibody, PGP-1 Antibody, Pgp1 Antibody, Phagocytic glycoprotein I Antibody, MDU3 Antibody, CD44R Antibody, HCELL Antibody, Heparan sulfate proteoglycan Antibody |
| Host | CD44 antibody was produced in Mouse |
| Clonality: | Monoclonal |
| Isotype: | IgG1 |
| Conjugations: | Fluorescein (FITC) |
| Immunogen Species: | CD44 antibody was raised against Bovine |
| Immunogen: | CD44 antibody was raised against bovine CD44. |
| Specificity: | Recognizes bovine CD44. Species cross-reactivity: sheep. |
| Reactivity: | Bovine, Sheep |
| Purification: | Protein G purified |
| Presentation: | PBS, pH 7.4, 1% BSA, 0.09% sodium azide |
| Recommended Storage: | Short term +4C. Aliquot to avoid freeze/thaw cycles and store at -20C. FITC conjugates are sensitive to light. |
| Usage Summary: | Suitable for use in Flow cytometry: Neat; 10 ul labels 10^6 cells in 100ul. |
| Uses: | Flow Cytometry (Optimal dilution to be determined by the researcher) |
| Size: | 100 µg |
| Concentration: | 0.1 mg/ml |
| Requested From: | Japan |
| La | boratory Reagent For In Vitro Research Use Only |
| Not for resale without prior written consent from LifeSpan BioSciences, Inc. | |
| Created on 9/24/2014 | |