

| TP103 AntibodyFamily / Subfamily:Protease / Serine S9BHostDPP4 antibody was produced in MouseClonality:MonoclonalIsotype:IgMConjugations:Phycoerythrin (PE)Immunogen Species:DPP4 / CD26 antibody was raised against HumanSpecificity:Recognizes human CD26.Reactivity:HumanPurification:PurifiedPresentation:PBS, 0.1% sodium azide, stabilizing agent.Recommended Storage:Long term: Add glycerol (40-50%) -20°C; Short term: +4°CUsage Summary:Flow cytometry. Immunoprecipitation. Immunohistochemistry (frozen sections).   | DPP4 Antibody, ADCP-2 Antibody, ADABP Antibody, Dipeptidyl peptidase IV     Antibody, Dipeptidyl-peptidase 4 Antibody, DPPIV Antibody, ADCP2 Antibody, T-     cell activation antigen CD26 Antibody, CD26 Antibody, CD26 antigen Antibody,     Dipeptidyl peptidase 4 Antibody, Dipeptidylpeptidase 4 Antibody, DPP IV Antibody, TP103 Antibody     Protease / Serine S9B     DPP4 antibody was produced in Mouse |
|--|---|
| Antibody, Dipeptidyl-peptidase 4 Antibody, DPPIV Antibody, ADCP2 Antibody, Dipeptidyl peptidase 4 Antibody, CD26 Antibody, CD26 Antibody, DPP IV Antibody, TP103 Antibody     Family / Subfamily:   Protease / Serine S9B     Host   DPP4 antibody was produced in Mouse     Clonality:   Monoclonal     Isotype:   IgM     Conjugations:   Phycoerythrin (PE)     Immunogen Species:   DPP4 / CD26 antibody was raised against Human     Specificity:   Recognizes human CD26.     Reactivity:   Human     Purification:   Purified     Presentation:   PBS, 0.1% sodium azide, stabilizing agent.     Recommended Storage:   Long term: Add glycerol (40-50%) -20°C; Short term: +4°C     Usage Summary:   Flow cytometry. Immunoprecipitation, Immunohistochemistry (frozen sections).     Studies of CD26 mediated T cell signaling. Studies of involvement of CD26 in HIV infection. Flow Cytometry: 10 ul/1E6 cells.     Uses:   IHC - Frozen, Immunoprecipitation, Flow Cytometry (Optimal dilution to be determined by the researcher)     Size:   100 tst     Requested From:   Japan     Laboratory Reagent For In Vitro Research Use Only | Antibody, Dipeptidyl-peptidase 4 Antibody, DPPIV Antibody, ADCP2 Antibody, T-<br>cell activation antigen CD26 Antibody, CD26 Antibody, CD26 antigen Antibody,<br>Dipeptidyl peptidase 4 Antibody, Dipeptidylpeptidase 4 Antibody, DPP IV Antibody,<br>TP103 Antibody<br>Protease / Serine S9B<br>DPP4 antibody was produced in Mouse<br>Monoclonal  |
| Host   DPP4 antibody was produced in Mouse     Clonality:   Monoclonal     Isotype:   IgM     Conjugations:   Phycoerythrin (PE)     Immunogen Species:   DPP4 / CD26 antibody was raised against Human     Specificity:   Recognizes human CD26.     Reactivity:   Human     Purification:   Purified     Presentation:   PBS, 0.1% sodium azide, stabilizing agent.     Recommended Storage:   Long term: Add glycerol (40-50%) -20°C; Short term: +4°C     Usage Summary:   Flow cytometry. Immunoprecipitation. Immunohistochemistry (frozen sections).<br>Studies of CD26 mediated T cell signaling. Studies of involvement of CD26 in HIV<br>infection. Flow Cytometry: 10 ul/1E6 cells.     Uses:   IHC - Frozen, Immunoprecipitation, Flow Cytometry (Optimal dilution to be<br>determined by the researcher)     Size:   100 tst     Requested From:   Japan     Laboratory Reagent For In Vitro Research Use Only  | DPP4 antibody was produced in Mouse<br>Monoclonal   |
| Clonality:   Monoclonal     Isotype:   IgM     Conjugations:   Phycoerythrin (PE)     Immunogen Species:   DPP4 / CD26 antibody was raised against Human     Specificity:   Recognizes human CD26.     Reactivity:   Human     Purification:   Purified     Presentation:   PBS, 0.1% sodium azide, stabilizing agent.     Recommended Storage:   Long term: Add glycerol (40-50%) -20°C; Short term: +4°C     Usage Summary:   Flow cytometry. Immunoprecipitation. Immunohistochemistry (frozen sections).<br>Studies of CD26 mediated T cell signaling. Studies of involvement of CD26 in HIV<br>infection. Flow Cytometry: 10 ul/1E6 cells.     Uses:   IHC - Frozen, Immunoprecipitation, Flow Cytometry (Optimal dilution to be<br>determined by the researcher)     Size:   100 tst     Requested From:   Japan     Laboratory Reagent For In Vitro Research Use Only   | Monoclonal  |
| Isotype:   IgM     Conjugations:   Phycoerythrin (PE)     Immunogen Species:   DPP4 / CD26 antibody was raised against Human     Specificity:   Recognizes human CD26.     Reactivity:   Human     Purification:   Purified     Presentation:   PBS, 0.1% sodium azide, stabilizing agent.     Recommended Storage:   Long term: Add glycerol (40-50%) -20°C; Short term: +4°C     Usage Summary:   Flow cytometry. Immunoprecipitation. Immunohistochemistry (frozen sections).<br>Studies of CD26 mediated T cell signaling. Studies of involvement of CD26 in HIV<br>infection. Flow Cytometry: 10 ul/1E6 cells.     Uses:   IHC - Frozen, Immunoprecipitation, Flow Cytometry (Optimal dilution to be<br>determined by the researcher)     Size:   100 tst     Requested From:   Japan     Laboratory Reagent For In Vitro Research Use Only   |   |
| Conjugations:   Phycoerythrin (PE)     Immunogen Species:   DPP4 / CD26 antibody was raised against Human     Specificity:   Recognizes human CD26.     Reactivity:   Human     Purification:   Purified     Presentation:   PBS, 0.1% sodium azide, stabilizing agent.     Recommended Storage:   Long term: Add glycerol (40-50%) -20°C; Short term: +4°C     Usage Summary:   Flow cytometry. Immunoprecipitation. Immunohistochemistry (frozen sections).<br>Studies of CD26 mediated T cell signaling. Studies of involvement of CD26 in HIV<br>infection. Flow Cytometry: 10 ul/1E6 cells.     Uses:   IHC - Frozen, Immunoprecipitation, Flow Cytometry (Optimal dilution to be<br>determined by the researcher)     Size:   100 tst     Requested From:   Japan     Laboratory Reagent For In Vitro Research Use Only  | IgM   |
| Immunogen Species:   DPP4 / CD26 antibody was raised against Human     Specificity:   Recognizes human CD26.     Reactivity:   Human     Purification:   Purified     Presentation:   PBS, 0.1% sodium azide, stabilizing agent.     Recommended Storage:   Long term: Add glycerol (40-50%) -20°C; Short term: +4°C     Usage Summary:   Flow cytometry. Immunoprecipitation. Immunohistochemistry (frozen sections).<br>Studies of CD26 mediated T cell signaling. Studies of involvement of CD26 in HIV<br>infection. Flow Cytometry: 10 ul/1E6 cells.     Uses:   IHC - Frozen, Immunoprecipitation, Flow Cytometry (Optimal dilution to be<br>determined by the researcher)     Size:   100 tst     Requested From:   Japan     Laboratory Reagent For In Vitro Research Use Only   |   |
| Specificity:   Recognizes human CD26.     Reactivity:   Human     Purification:   Purified     Presentation:   PBS, 0.1% sodium azide, stabilizing agent.     Recommended Storage:   Long term: Add glycerol (40-50%) -20°C; Short term: +4°C     Usage Summary:   Flow cytometry. Immunoprecipitation. Immunohistochemistry (frozen sections).<br>Studies of CD26 mediated T cell signaling. Studies of involvement of CD26 in HIV<br>infection. Flow Cytometry: 10 ul/1E6 cells.     Uses:   IHC - Frozen, Immunoprecipitation, Flow Cytometry (Optimal dilution to be<br>determined by the researcher)     Size:   100 tst     Requested From:   Japan     Laboratory Reagent For In Vitro Research Use Only  | Phycoerythrin (PE)  |
| Reactivity:   Human     Purification:   Purified     Presentation:   PBS, 0.1% sodium azide, stabilizing agent.     Recommended Storage:   Long term: Add glycerol (40-50%) -20°C; Short term: +4°C     Usage Summary:   Flow cytometry. Immunoprecipitation. Immunohistochemistry (frozen sections).<br>Studies of CD26 mediated T cell signaling. Studies of involvement of CD26 in HIV<br>infection. Flow Cytometry: 10 ul/1E6 cells.     Uses:   IHC - Frozen, Immunoprecipitation, Flow Cytometry (Optimal dilution to be<br>determined by the researcher)     Size:   100 tst     Requested From:   Japan     Laboratory Reagent For In Vitro Research Use Only  | DPP4 / CD26 antibody was raised against Human   |
| Purification:   Purified     Presentation:   PBS, 0.1% sodium azide, stabilizing agent.     Recommended Storage:   Long term: Add glycerol (40-50%) -20°C; Short term: +4°C     Usage Summary:   Flow cytometry. Immunoprecipitation. Immunohistochemistry (frozen sections).<br>Studies of CD26 mediated T cell signaling. Studies of involvement of CD26 in HIV<br>infection. Flow Cytometry: 10 ul/1E6 cells.     Uses:   IHC - Frozen, Immunoprecipitation, Flow Cytometry (Optimal dilution to be<br>determined by the researcher)     Size:   100 tst     Requested From:   Japan     Laboratory Reagent For In Vitro Research Use Only  | Recognizes human CD26.  |
| Presentation:   PBS, 0.1% sodium azide, stabilizing agent.     Recommended Storage:   Long term: Add glycerol (40-50%) -20°C; Short term: +4°C     Usage Summary:   Flow cytometry. Immunoprecipitation. Immunohistochemistry (frozen sections).<br>Studies of CD26 mediated T cell signaling. Studies of involvement of CD26 in HIV<br>infection. Flow Cytometry: 10 ul/1E6 cells.     Uses:   IHC - Frozen, Immunoprecipitation, Flow Cytometry (Optimal dilution to be<br>determined by the researcher)     Size:   100 tst     Requested From:   Japan     Laboratory Reagent For In Vitro Research Use Only   | Human   |
| Recommended Storage:   Long term: Add glycerol (40-50%) -20°C; Short term: +4°C     Usage Summary:   Flow cytometry. Immunoprecipitation. Immunohistochemistry (frozen sections).<br>Studies of CD26 mediated T cell signaling. Studies of involvement of CD26 in HIV<br>infection. Flow Cytometry: 10 ul/1E6 cells.     Uses:   IHC - Frozen, Immunoprecipitation, Flow Cytometry (Optimal dilution to be<br>determined by the researcher)     Size:   100 tst     Requested From:   Japan     Laboratory Reagent For In Vitro Research Use Only  | Purified  |
| Usage Summary:   Flow cytometry. Immunoprecipitation. Immunohistochemistry (frozen sections).<br>Studies of CD26 mediated T cell signaling. Studies of involvement of CD26 in HIV<br>infection. Flow Cytometry: 10 ul/1E6 cells.     Uses:   IHC - Frozen, Immunoprecipitation, Flow Cytometry (Optimal dilution to be<br>determined by the researcher)     Size:   100 tst     Requested From:   Japan     Laboratory Reagent For In Vitro Research Use Only  | PBS, 0.1% sodium azide, stabilizing agent.  |
| Studies of CD26 mediated T cell signaling. Studies of involvement of CD26 in HIV infection. Flow Cytometry: 10 ul/1E6 cells.     Uses:   IHC - Frozen, Immunoprecipitation, Flow Cytometry (Optimal dilution to be determined by the researcher)     Size:   100 tst     Requested From:   Japan     Laboratory Reagent For In Vitro Research Use Only   | Long term: Add glycerol (40-50%) -20°C; Short term: +4°C  |
| determined by the researcher)   Size:   100 tst   Requested From:   Japan   Laboratory Reagent For In Vitro Research Use Only  | Studies of CD26 mediated T cell signaling. Studies of involvement of CD26 in HIV  |
| Requested From:   Japan     Laboratory Reagent For In Vitro Research Use Only  | IHC - Frozen, Immunoprecipitation, Flow Cytometry (Optimal dilution to be determined by the researcher)   |
| Laboratory Reagent For In Vitro Research Use Only  | 100 tst   |
|  | Japan   |
| Not for resale without prior written consent from LifeSpan BioSciences, Inc.   | oratory Reagent For In Vitro Research Use Only  |
| Created on 9/25/2014   |   |
|  |   |