


**NMI Mouse anti-Human Monoclonal (9E8) Antibody - LS-C133104 - LSBio**


<b>CatalogID:</b>	LS-C133104
<b>Target:</b>	N-myc (and STAT) interactor (NMI)
<b>Synonyms:</b>	NMI Antibody, Hou Antibody, N-myc and STAT interactor Antibody, N-myc-interactor Antibody, N-myc (and STAT) interactor Antibody, N-myc interactor Antibody
<b>Host</b>	NMI antibody was produced in Mouse
<b>Clonality:</b>	Monoclonal
<b>Isotype:</b>	IgG2b,k
<b>Clone Name:</b>	9E8
<b>Immunogen Species:</b>	NMI antibody was raised against Human
<b>Immunogen:</b>	NMI antibody was raised against nMI (AAH01268, 1 a.a. ~ 101 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
<b>Specificity:</b>	Human Hou
<b>Reactivity:</b>	Human
<b>Purification:</b>	Protein A purified
<b>Presentation:</b>	PBS, pH 7.2
<b>Recommended Storage:</b>	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
<b>Usage Summary:</b>	Immunohistochemistry: Formalin-fixed paraffin-embedded sections. Sandwich ELISA: Recombinant protein. Western Blot: Cell lysate, Recombinant protein.
<b>Uses:</b>	IHC - Paraffin (3 µg/ml), Western blot, ELISA (Optimal dilution to be determined by the researcher)
<b>Size:</b>	100 µg

**Immunohistochemistry Image:**




Immunoperoxidase of monoclonal antibody to NMI on formalin-fixed paraffin-embedded human small Intestine. [antibody concentration 3 ug/ml]

**Western Blot Image:**



NMI monoclonal antibody (M04), clone 9E8 Western blot of NMI expression in HeLa.

**ELISA Image:**



Detection limit for recombinant GST tagged NMI is approximately 0.1 ng/ml as a capture antibody.

**Requested From:**

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/25/2014

© 2014 LifeSpan BioSciences