

**MYC / c-Myc Sheep anti-Human Polyclonal Antibody - LS-C136227 - LSBio**

<b>CatalogID:</b>	LS-C136227
<b>Target:</b>	v-myc avian myelocytomatosis viral oncogene homolog
<b>Synonyms:</b>	MYC Antibody, BHLHe39 Antibody, C-Myc Antibody, MRTL Antibody, Myc proto-oncogene protein Antibody, Proto-oncogene c-Myc Antibody, Rats1 Antibody, Transcription factor p64 Antibody, v-myc Antibody
<b>Host</b>	MYC antibody was produced in Sheep
<b>Clonality:</b>	Polyclonal
<b>Immunogen Species:</b>	MYC / c-Myc antibody was raised against Human
<b>Antigen Type:</b>	Synthetic peptide
<b>Immunogen:</b>	MYC / c-Myc antibody was raised against synthetic peptide {EQKLISEEDL (410-419)} from human c-Myc (c Myc tag) conjugated to an immunogenic carrier protein was used as the antigen.
<b>Specificity:</b>	Specific for c-Myc.
<b>Reactivity:</b>	Human
<b>Purification:</b>	Purified
<b>Reconstitution:</b>	Sterile water. Possible additional volumes for resuspension: 500 µl
<b>Presentation:</b>	Lyophilized. Centrifuge to remove any insoluble material
<b>Recommended Storage:</b>	Maintain the lyophilized/reconstituted antibodies frozen at -20°C for long term storage and refrigerated at 2-8°C for a shorter term. When reconstituting, glycerol (1:1) may be added for an additional stability. Avoid freeze and thaw cycles.
<b>Uses:</b>	Western blot (10 - 50 µg/ml) (Optimal dilution to be determined by the researcher)
<b>Size:</b>	500 µg
<b>Concentration:</b>	1 µg/µl (after resuspension)
<b>Requested From:</b>	Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/25/2014

© 2014 LifeSpan BioSciences