

IP-10 / CXCL10 Mouse anti-Human Monoclonal (FITC) (6D4) Antibody - LS-C139962 - LSBio	
<b>CatalogID:</b>	LS-C139962
<b>Target:</b>	chemokine (C-X-C motif) ligand 10 (CXCL10)
<b>Synonyms:</b>	CXCL10 Antibody, C-X-C motif chemokine 10 Antibody, Gamma-IP10 Antibody, IP-10 Antibody, Gamma IP10 Antibody, GIP-10 Antibody, Mob-1 Antibody, Small-inducible cytokine B10 Antibody, Crg-2 Antibody, IFI10 Antibody, INP10 Antibody, SCYB10 Antibody
<b>Family / Subfamily:</b>	Intercrine / Intercrine alpha
<b>Host</b>	CXCL10 antibody was produced in Mouse
<b>Clonality:</b>	Monoclonal
<b>Isotype:</b>	IgG2a
<b>Clone Name:</b>	6D4
<b>Conjugations:</b>	Fluorescein (FITC)
<b>Immunogen Species:</b>	IP-10 / CXCL10 antibody was raised against Human
<b>Immunogen:</b>	IP-10 / CXCL10 antibody was raised against recombinant IP-10 (human)
<b>Specificity:</b>	The monoclonal antibody 6D4 recognizes human C-X-C motif chemokine 10 (IP-10), a protein of 98 amino acids.
<b>Reactivity:</b>	Human
<b>Purification:</b>	Purified
<b>Presentation:</b>	PBS, 0.1% BSA, 0.02% sodium azide
<b>Recommended Storage:</b>	Store at 4°C, stable for one year.
<b>Usage Summary:</b>	For immunohistochemistry, flow cytometry and Western blotting, dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50.
<b>Uses:</b>	IHC - Frozen, Immunofluorescence, Western blot, Flow Cytometry (Optimal dilution to be determined by the researcher)
<b>Size:</b>	100 µg
<b>Requested From:</b>	Japan
Laboratory Reagent For In Vitro Research Use Only	
Not for resale without prior written consent from LifeSpan BioSciences, Inc.	
Created on 9/25/2014	
© 2014 LifeSpan BioSciences	