

Antibody, IIAE2 Antibody Family / Subfamily: Toll-like Receptor / not assigned-Toll-like Receptor Host TLR3 antibody was produced in Mouse Clonality: Monoclonal Isotype: IgG1 Clone Name: TLR3.7 Conjugations: Fluorescein (FITC) Immunogen Species: TLR3 antibody was raised against Human Immunogen: TLR3 antibody was raised against human Flag-tagged TLR3 stably expressed by Ba/F3 cells Specificity: Human TLR3 Reactivity: Human, Mouse, Dog Purification: Purified Presentation: PBS, 0.1% BSA Recommended Storage: Store at 4°C, stable for one year. Usage Summary: For flow cytometry dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical stating working dilution is 1:50. Uses: IHC - Paraffin, IHC - Frozen, Immunofluorescence, Western blot, Immunoprecipitation, Flow Cytometry, Functional Assay (Optimal dilution to be determined by the researcher) Size: 100 µg Concentration: 0.1 mg/ml	CatalogID:	LS-C140026
Antibody, IIAE2 Antibody Family / Subfamily: Toll-like Receptor / not assigned-Toll-like Receptor Host TLR3 antibody was produced in Mouse Clonality: Monoclonal Isotype: IgG1 Clone Name: TLR3.7 Conjugations: Fluorescein (FITC) Immunogen Species: TLR3 antibody was raised against Human Immunogen: TLR3 antibody was raised against human Flag-tagged TLR3 stably expressed by Ba/F3 cells Specificity: Human TLR3 Reactivity: Human, Mouse, Dog Purification: Purified Presentation: PBS, 0.1% BSA Recommended Storage: Store at 4°C, stable for one year. Usage Summary: For flow cytometry dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50. Uses: IHC - Paraffin, IHC - Forzen, Immunofluorescence, Western blot, Immunoprecipitaton, Flow Cytometry, Functional Assay (Optimal dilution to be determined by the researcher) Size: 100 µg Concentration: 0.1 mg/ml Requested From: Japan	Target:	toll-like receptor 3 (TLR3)
Host TLR3 antibody was produced in Mouse Clonality: Monoclonal Isotype: IgG1 Clone Name: TLR3.7 Conjugations: Fluorescein (FITC) Immunogen Species: TLR3 antibody was raised against Human Immunogen: TLR3 antibody was raised against Human Immunogen: TLR3 antibody was raised against human Flag-tagged TLR3 stably expressed by Ba/F3 cells Specificity: Human TLR3 Reactivity: Human, Mouse, Dog Purification: Pusified Presentation: PBS, 0.1% BSA Recommended Storage: Store at 4°C, stable for one year. Usage Summary: For flow cytometry dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. Tool. Uses: IHC - Paraffin, IHC - Frozen, Immunofluorescence, Western blot, Immunoprecipitation, Flow Cytometry, Functional Assay (Optimal dilution to be determined by the researcher) Size: 100 µg Concentration: 0.1 mg/ml Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	Synonyms:	TLR3 Antibody, CD283 Antibody, Toll-like receptor 3 Antibody, CD283 antigen Antibody, IIAE2 Antibody
Clonality: Monoclonal Isotype: IgG1 Clone Name: TLR3.7 Conjugations: Fluorescein (FITC) Immunogen Species: TLR3 antibody was raised against Human Immunogen: TLR3 antibody was raised against human Flag-tagged TLR3 stably expressed by Ba/F3 cells Specificity: Human TLR3 Reactivity: Human, Mouse, Dog Purification: Purified Presentation: PBS, 0.1% BSA Recommended Storage: Store at 4°C, stable for one year. Usage Summary: For flow cytometry dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50. Uses: IHC - Paraffin, IHC - Frozen, Immunofluorescence, Western blot, Immunoprecipitation, Flow Cytometry, Functional Assay (Optimal dilution to be determined by the researcher) Size: 100 µg Concentration: 0.1 mg/ml Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	Family / Subfamily:	Toll-like Receptor / not assigned-Toll-like Receptor
Isotype: IgG1 Clone Name: TLR3.7 Conjugations: Fluorescein (FITC) Immunogen Species: TLR3 antibody was raised against Human Immunogen: TLR3 antibody was raised against human Flag-tagged TLR3 stably expressed by Ba/F3 cells Specificity: Human TLR3 Reactivity: Human, Mouse, Dog Purification: Purified Presentation: PBS, 0.1% BSA Recommended Storage: Store at 4°C, stable for one year. Usage Summary: For flow cytometry dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50. Uses: IHC - Paraffin, IHC - Frozen, Immunofluorescence, Western blot, Immunoprecipitation, Flow Cytometry, Functional Assay (Optimal dilution to be determined by the researcher) Size: 100 µg Concentration: 0.1 mg/ml Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	Host	TLR3 antibody was produced in Mouse
Clone Name: TLR3.7 Conjugations: Fluorescein (FITC) Immunogen Species: TLR3 antibody was raised against Human Immunogen: TLR3 antibody was raised against human Flag-tagged TLR3 stably expressed by Ba/F3 cells Specificity: Human TLR3 Reactivity: Human, Mouse, Dog Purification: Purified Presentation: PBS, 0.1% BSA Recommended Storage: Store at 4°C, stable for one year. Usage Summary: For flow cytometry dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50. Uses: IHC - Paraffin, IHC - Frozen, Immunofluorescence, Western blot, Immunoprecipitation, Flow Cytometry, Functional Assay (Optimal dilution to be determined by the researcher) Size: 100 µg Concentration: 0.1 mg/ml Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	Clonality:	Monoclonal
Conjugations: Fluorescein (FITC) Immunogen Species: TLR3 antibody was raised against Human Immunogen: TLR3 antibody was raised against human Flag-tagged TLR3 stably expressed by Ba/F3 cells Specificity: Human TLR3 Reactivity: Human, Mouse, Dog Purification: Purified Presentation: PBS, 0.1% BSA Recommended Storage: Store at 4°C, stable for one year. Usage Summary: For flow cytometry dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50. Uses: IHC - Paraffin, IHC - Frozen, Immunofluorescence, Western blot, Immunoprecipitation, Flow Cytometry, Functional Assay (Optimal dilution to be determined by the researcher) Size: 100 μg Concentration: 0.1 mg/ml Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	Isotype:	lgG1
Immunogen Species: TLR3 antibody was raised against Human Immunogen: TLR3 antibody was raised against human Flag-tagged TLR3 stably expressed by Ba/F3 cells Specificity: Human TLR3 Reactivity: Human, Mouse, Dog Purification: Purified Presentation: PBS, 0.1% BSA Recommended Storage: Store at 4°C, stable for one year. Usage Summary: For flow cytometry dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50. Uses: IHC - Paraffin, IHC - Frozen, Immunofluorescence, Western blot, Immunoprecipitation, Flow Cytometry, Functional Assay (Optimal dilution to be determined by the researcher) Size: 100 µg Concentration: 0.1 mg/ml Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	Clone Name:	TLR3.7
Immunogen: TLR3 antibody was raised against human Flag-tagged TLR3 stably expressed by Ba/F3 cells Specificity: Human TLR3 Reactivity: Human, Mouse, Dog Purification: Purified Presentation: PBS, 0.1% BSA Recommended Storage: Store at 4°C, stable for one year. Usage Summary: For flow cytometry dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50. Uses: IHC - Paraffin, IHC - Frozen, Immunofluorescence, Western blot, Immunoprecipitation, Flow Cytometry, Functional Assay (Optimal dilution to be determined by the researcher) Size: 100 µg Concentration: 0.1 mg/ml Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	Conjugations:	Fluorescein (FITC)
Ba/F3 cells Ba/F3 cells Specificity: Human TLR3 Reactivity: Human, Mouse, Dog Purification: Purified Presentation: PBS, 0.1% BSA Recommended Storage: Store at 4°C, stable for one year. Usage Summary: For flow cytometry dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50. Uses: IHC - Paraffin, IHC - Frozen, Immunofluorescence, Western blot, Immunoprecipitation, Flow Cytometry, Functional Assay (Optimal dilution to be determined by the researcher) Size: 100 µg Concentration: 0.1 mg/ml Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	Immunogen Species:	TLR3 antibody was raised against Human
Reactivity: Human, Mouse, Dog Purification: Purified Presentation: PBS, 0.1% BSA Recommended Storage: Store at 4°C, stable for one year. Usage Summary: For flow cytometry dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50. Uses: IHC - Paraffin, IHC - Frozen, Immunofluorescence, Western blot, Immunoprecipitation, Flow Cytometry, Functional Assay (Optimal dilution to be determined by the researcher) Size: 100 µg Concentration: 0.1 mg/ml Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	Immunogen:	TLR3 antibody was raised against human Flag-tagged TLR3 stably expressed by Ba/F3 cells
Purification: Purified Presentation: PBS, 0.1% BSA Recommended Storage: Store at 4°C, stable for one year. Usage Summary: For flow cytometry dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50. Uses: IHC - Paraffin, IHC - Frozen, Immunofluorescence, Western blot, Immunoprecipitation, Flow Cytometry, Functional Assay (Optimal dilution to be determined by the researcher) Size: 100 μg Concentration: 0.1 mg/ml Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	Specificity:	Human TLR3
Presentation: PBS, 0.1% BSA Recommended Storage: Store at 4°C, stable for one year. Usage Summary: For flow cytometry dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50. Uses: IHC - Paraffin, IHC - Frozen, Immunofluorescence, Western blot, Immunoprecipitation, Flow Cytometry, Functional Assay (Optimal dilution to be determined by the researcher) Size: 100 µg Concentration: 0.1 mg/ml Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	Reactivity:	Human, Mouse, Dog
Recommended Storage: Store at 4°C, stable for one year. Usage Summary: For flow cytometry dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50. Uses: IHC - Paraffin, IHC - Frozen, Immunofluorescence, Western blot, Immunoprecipitation, Flow Cytometry, Functional Assay (Optimal dilution to be determined by the researcher) Size: 100 μg Concentration: 0.1 mg/ml Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	Purification:	Purified
Usage Summary: For flow cytometry dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50. Uses: IHC - Paraffin, IHC - Frozen, Immunofluorescence, Western blot, Immunoprecipitation, Flow Cytometry, Functional Assay (Optimal dilution to be determined by the researcher) Size: 100 μg Concentration: 0.1 mg/ml Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	Presentation:	PBS, 0.1% BSA
recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50. Uses: IHC - Paraffin, IHC - Frozen, Immunofluorescence, Western blot, Immunoprecipitation, Flow Cytometry, Functional Assay (Optimal dilution to be determined by the researcher) Size: 100 μg Concentration: 0.1 mg/ml Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	Recommended Storage:	Store at 4°C, stable for one year.
Immunoprecipitation, Flow Cytometry, Functional Assay (Optimal dilution to be determined by the researcher) Size: 100 μg Concentration: 0.1 mg/ml Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	Usage Summary:	recommended that users test the reagent and determine their own optimal
Concentration: 0.1 mg/ml Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	Uses:	Immunoprecipitation, Flow Cytometry, Functional Assay (Optimal dilution to be
Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	Size:	100 µg
Laboratory Reagent For In Vitro Research Use Only	Concentration:	0.1 mg/ml
	Requested From:	Japan
Not for resale without prior written consent from LifeSpan BioSciences, Inc.	Lal	poratory Reagent For In Vitro Research Use Only
Created on 9/25/2014	Not for resale	