

Proteinase 3 / PRTN3 Mouse anti-Human Monoclonal (FITC) (PR3-G2) Antibody - LS-C140094 - LSBio	
<b>CatalogID:</b>	LS-C140094
<b>Target:</b>	proteinase 3 (PRTN3)
<b>Synonyms:</b>	PRTN3 Antibody, ACPA Antibody, Azurophil granule protein 7 Antibody, C-ANCA antigen Antibody, CANCA Antibody, C-ANCA Antibody, AGP7 Antibody, Leukocyte proteinase 3 Antibody, MBN Antibody, Myeloblastin Antibody, NP-4 Antibody, Neutrophil proteinase 4 Antibody, NP4 Antibody, p29 Antibody, PR-3 Antibody, Proteinase 3 Antibody, MBT Antibody, Wegener autoantigen Antibody, PR3 Antibody, Serine proteinase, neutrophil Antibody
<b>Family / Subfamily:</b>	Protease / Serine S1
<b>Host</b>	PRTN3 antibody was produced in Mouse
<b>Clonality:</b>	Monoclonal
<b>Isotype:</b>	IgG1
<b>Clone Name:</b>	PR3-G2
<b>Conjugations:</b>	Fluorescein (FITC)
<b>Immunogen Species:</b>	Proteinase 3 / PRTN3 antibody was raised against Human
<b>Specificity:</b>	Human PRTN3
<b>Reactivity:</b>	Human
<b>Purification:</b>	Purified
<b>Presentation:</b>	PBS, 1% BSA, 0.02% sodium azide
<b>Recommended Storage:</b>	Store at 4°C, stable for one year.
<b>Usage Summary:</b>	For immunohistology, flow cytometry and Western blotting dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50.
<b>Uses:</b>	IHC - Frozen, Western blot, Flow Cytometry (Optimal dilution to be determined by the researcher)
<b>Size:</b>	100 µg
<b>Concentration:</b>	0.1 mg/ml
<b>Requested From:</b>	Japan
<b>Laboratory Reagent For In Vitro Research Use Only</b> Not for resale without prior written consent from LifeSpan BioSciences, Inc. Created on 9/25/2014 © 2014 LifeSpan BioSciences	