

| MME / CD10 Mouse anti-Human Monoclonal (FITC) (RXB.14) Antibody - LS-C140284 - LSBio |   |
|--|---|
| CatalogID:   | LS-C140284  |
| Target:  | membrane metallo-endopeptidase (MME)  |
| Synonyms:  | MME Antibody, Atriopeptidase Antibody, CALLA Antibody, CD10 Antibody, EPN Antibody, Membrane metallo-endopeptidase Antibody, NEP Antibody, Neprilysin Antibody, Neutral endopeptidase Antibody, Neutral endopeptidase 24.11 Antibody, CD10 antigen Antibody, Enkephalinase Antibody, Neprylisin Antibody, SFE Antibody, Skin fibroblast elastase Antibody   |
| Family / Subfamily:  | Protease / Metallopeptidase M13   |
| Host   | MME antibody was produced in Mouse  |
| Clonality:   | Monoclonal  |
| Isotype:   | IgG1  |
| Clone Name:  | RXB.14  |
| Conjugations:  | Fluorescein (FITC)  |
| Immunogen Species:   | MME / CD10 antibody was raised against Human  |
| Specificity:   | Human MME / CD10  |
| Reactivity:  | Human   |
| Purification:  | Protein A/G purified  |
| Presentation:  | PBS, 0.08% sodium azide, 0.2% carrier protein, sterile-filtered   |
| Recommended Storage:   | Store at 4°C. Do not freeze.  |
| Usage Summary:   | PBMC: Add 10 ul of antibody/10^6 PBMC in 100 ul PBS. Mix gently and incubate for 15 minutes at 2 to 8°C. Wash twice with PBS and analyze. WHOLE BLOOD: Add 10 ul of antibody/100 ul of whole blood. Mix gently and incubate for 15 minutes at room temperature 20°C. Lyse the whole blood. Wash once with PBS and analyze. See instrument manufacturers instructions for Lysed Whole Blood and Immunofluorescence analysis with a flow cytometer or microscope. |
| Uses:  | Flow Cytometry (Optimal dilution to be determined by the researcher)  |
| Size:  | 100 tst   |
| Requested From:  | Japan   |

Laboratory Reagent For In Vitro Research Use Only
Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/25/2014 © 2014 LifeSpan BioSciences