

	LS-C140319
Farget:	interleukin 2 receptor, alpha (IL2RA)
Synonyms:	IL2RA Antibody, CD25 antigen Antibody, CD25 Antibody, IDDM10 Antibody, IL-2 receptor subunit alpha Antibody, IL2R Antibody, Interleukin 2 receptor, alpha Antibody, IL-2-RA Antibody, IL2-RA Antibody, TCGFR Antibody, IL-2R subunit alpha Antibody, p55 Antibody, TAC antigen Antibody
Family / Subfamily:	Interleukin
lost	IL2RA antibody was produced in Mouse
Clonality:	Monoclonal
sotype:	lgG1
Clone Name:	ITYV
Conjugations:	Phycoerythrin (PE)
mmunogen Species:	IL2RA / CD25 antibody was raised against Human
mmunogen:	IL2RA / CD25 antibody was raised against pHA-activated human T-lymphocytes.
Specificity:	Human IL2RA / CD25
Reactivity:	Human
Purification:	Protein A/G purified
Presentation:	PBS, 0.08% sodium azide, 0.2% carrier protein, sterile-filtered
Recommended Storage:	Store at 4°C. Do not freeze.
Jsage Summary:	PBMC: Add 10 ul of antibody/10^6 PBMC in 100 ul PBS. Mix gently and incubate for 15 minutes at 2 to 8°C. Wash twice with PBS and analyze or fix with 0.5% v/v or paraformaldehyde in PBS and analyze. WHOLE BLOOD: Add 10 ul of antibody/100 ul of whole blood. Mix gently and incubate for 15 minutes at room temperature 20°C. Lyse the whole blood. Wash once with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze. See instrument manufacturers instructions for Lysed Whole Blood and Immunofluorescence analysis with a flow cytometer or microscope.
Jses:	Flow Cytometry (Optimal dilution to be determined by the researcher)
Size:	100 tst
Requested From:	Japan
La	aboratory Reagent For In Vitro Research Use Only
Not for resale	without prior written consent from LifeSpan BioSciences, Inc. Created on 9/25/2014