

DPP4 / CD26 Mouse anti-Human Monoclonal (FITC) (EUG-6) Antibody - LS-C140321 - LSBio	
<b>CatalogID:</b>	LS-C140321
<b>Target:</b>	dipeptidyl-peptidase 4 (DPP4)
<b>Synonyms:</b>	DPP4 Antibody, ADCP-2 Antibody, ADABP Antibody, Dipeptidyl peptidase IV Antibody, Dipeptidyl-peptidase 4 Antibody, DPPIV Antibody, ADCP2 Antibody, T-cell activation antigen CD26 Antibody, CD26 Antibody, CD26 antigen Antibody, Dipeptidyl peptidase 4 Antibody, Dipeptidylpeptidase 4 Antibody, DPP IV Antibody, TP103 Antibody
<b>Family / Subfamily:</b>	Protease / Serine S9B
<b>Host</b>	DPP4 antibody was produced in Mouse
<b>Clonality:</b>	Monoclonal
<b>Isotype:</b>	IgG1
<b>Clone Name:</b>	EUG-6
<b>Conjugations:</b>	Fluorescein (FITC)
<b>Immunogen Species:</b>	DPP4 / CD26 antibody was raised against Human
<b>Specificity:</b>	Human DPP4 / CD26
<b>Reactivity:</b>	Human
<b>Purification:</b>	Protein A/G purified
<b>Presentation:</b>	PBS, 0.08% sodium azide, 0.2% carrier protein, sterile-filtered
<b>Recommended Storage:</b>	Store at 4°C. Do not freeze.
<b>Usage Summary:</b>	PBMC: Add 10 ul of antibody/10 <sup>6</sup> PBMC in 100 ul PBS. Mix gently and incubate for 15 minutes at 2 to 8°C. Wash twice with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze. WHOLE BLOOD: Add 10 ul of antibody/100 ul of whole blood. Mix gently and incubate for 15 minutes at room temperature (20°C). Lyse the whole blood. Wash once with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze. See instrument manufacturers instructions for Lysed Whole Blood and Immunofluorescence analysis with a flow cytometer or microscope.
<b>Uses:</b>	Flow Cytometry (Optimal dilution to be determined by the researcher)
<b>Size:</b>	100 tst
<b>Requested From:</b>	Japan
Laboratory Reagent For In Vitro Research Use Only	
Not for resale without prior written consent from LifeSpan BioSciences, Inc.	
Created on 9/25/2014	
© 2014 LifeSpan BioSciences	