

CD27 Mouse anti-Human Monoclonal (FITC) (DEN-3) Antibody - LS-C140324 - LSBio

CatalogID:	LS-C140324
Target:	CD27 molecule
Synonyms:	CD27 Antibody, CD27 antigen Antibody, CD27 molecule Antibody, S152 Antibody, T14 Antibody, T-cell activation antigen CD27 Antibody, CD27L receptor Antibody, T cell activation antigen S152 Antibody, TNFRSF7 Antibody, Tp55 Antibody
Family / Subfamily:	TNF Receptor / not assigned-TNF Receptor
Host	CD27 antibody was produced in Mouse
Clonality:	Monoclonal
Isotype:	IgG1
Clone Name:	DEN-3
Conjugations:	Fluorescein (FITC)
Immunogen Species:	CD27 antibody was raised against Human
Specificity:	Human CD27
Reactivity:	Human
Purification:	Protein A/G purified
Presentation:	PBS, 0.08% sodium azide, 0.2% carrier protein, sterile-filtered
Recommended Storage:	Store at 4°C. Do not freeze.
Usage Summary:	PBMC: Add 10 ul of antibody/10 ⁶ PBMC in 100 ul PBS. Mix gently and incubate for 15 minutes at 2 to 8°C. Wash twice with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze. WHOLE BLOOD: Add 10 ul of antibody/100 ul of whole blood. Mix gently and incubate for 15 minutes at room temperature (20°C). Lyse the whole blood. Wash once with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze. See instrument manufacturers instructions for Lysed Whole Blood and Immunofluorescence analysis with a flow cytometer or microscope.
Uses:	Flow Cytometry (Optimal dilution to be determined by the researcher)
Size:	100 tst
Requested From:	Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/25/2014

© 2014 LifeSpan BioSciences