

NCAM / CD56 Mouse anti-Human Monoclonal (FITC) (C5.9) Antibody - LS-C140369 - LSBio	
CatalogID:	LS-C140369
Target:	neural cell adhesion molecule 1 (NCAM1)
Synonyms:	NCAM1 Antibody, CD56 Antibody, CD56 antigen Antibody, MSK39 Antibody, NCAM-1 Antibody, N-CAM-1 Antibody, NCAM Antibody
Family / Subfamily:	Immunoglobulin / not assigned-Immunoglobulin
Host	NCAM1 antibody was produced in Mouse
Clonality:	Monoclonal
Isotype:	IgG2b
Clone Name:	C5.9
Conjugations:	Fluorescein (FITC)
Immunogen Species:	NCAM / CD56 antibody was raised against Human
Immunogen:	NCAM / CD56 antibody was raised against kG1a cell line.
Specificity:	Human NCAM1 / CD56
Reactivity:	Human
Presentation:	PBS, 0.08% sodium azide, 0.2% carrier protein, sterile-filtered
Recommended Storage:	Store at 4°C. Do not freeze.
Usage Summary:	PBMC: Add 10 ul of antibody/10 ⁶ PBMC in 100 ul PBS. Mix gently and incubate for 15 minutes at 2 to 8°C. Wash twice with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze. WHOLE BLOOD: Add 10 ul of antibody/100 ul of whole blood. Mix gently and incubate for 15 minutes at room temperature 20°C. Lyse the whole blood. Wash once with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze. See instrument manufacturers instructions for Lysed Whole Blood and Immunofluorescence analysis with a flow cytometer or microscope.
Uses:	Flow Cytometry (Optimal dilution to be determined by the researcher)
Size:	100 tst
Requested From:	Japan
Laboratory Reagent For In Vitro Research Use Only	
Not for resale without prior written consent from LifeSpan BioSciences, Inc.	
Created on 9/25/2014	
© 2014 LifeSpan BioSciences	