

SELL / L-Selectin / CD62L Mouse anti-Human Monoclonal (PE) (1H3) Antibody - LS-C140377 - LSBio	
CatalogID:	LS-C140377
Target:	selectin L (SELL)
Synonyms:	SELL Antibody, CD62L Antibody, CD62L antigen Antibody, LAM-1 Antibody, LAM1 Antibody, LECAM1 Antibody, LEU8 Antibody, Leukocyte adhesion molecule 1 Antibody, Lyam-1 Antibody, LSEL Antibody, gp90-MEL Antibody, HLHRc Antibody, PLNHR Antibody, L-selectin Antibody, Leu-8 Antibody, Selectin L Antibody, LNHR Antibody, LYAM1 Antibody, Lymph node homing receptor Antibody, Lymphocyte adhesion molecule 1 Antibody, TQ1 Antibody, Pln homing receptor Antibody
Host	SELL antibody was produced in Mouse
Clonality:	Monoclonal
Isotype:	IgG2a
Clone Name:	1H3
Conjugations:	Phycoerythrin (PE)
Immunogen Species:	SELL / L-Selectin / CD62L antibody was raised against Human
Immunogen:	SELL / L-Selectin / CD62L antibody was raised against peripheral blood T lymphocytes.
Specificity:	Human SELL
Reactivity:	Human
Purification:	Protein A/G purified
Presentation:	PBS, 0.08% sodium azide, 0.2% carrier protein, sterile-filtered
Recommended Storage:	Store at 4°C. Do not freeze.
Usage Summary:	PBMC: Add 10 ul of antibody/10^6 PBMC in 100 ul PBS. Mix gently and incubate for 15 minutes at 2 to 8°C. Wash twice with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze. WHOLE BLOOD: Add 10 ul of antibody/100 ul of whole blood. Mix gently and incubate for 15 minutes at room temperature (20°C). Lyse the whole blood. Wash once with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze. See instrument manufacturers instructions for Lysed Whole Blood and Immunofluorescence analysis with a flow cytometer or microscope.
Uses:	Flow Cytometry (Optimal dilution to be determined by the researcher)
Size:	100 tst
Requested From:	Japan

Laboratory Reagent For In Vitro Research Use Only
Not for resale without prior written consent from LifeSpan BioSciences, Inc.
Created on 9/25/2014

© 2014 LifeSpan BioSciences