

MCAM / CD146 Mouse anti-Human Monoclonal (FITC) Antibody - LS-C14276 - LSBio

<b>CatalogID:</b>	LS-C14276
<b>Target:</b>	melanoma cell adhesion molecule (MCAM)
<b>Synonyms:</b>	MCAM Antibody, CD146 Antibody, CD146 antigen Antibody, Gicerin Antibody, Melanoma adhesion molecule Antibody, MUC18 Antibody, p1h12 Antibody
<b>Family / Subfamily:</b>	Immunoglobulin / not assigned-Immunoglobulin
<b>Host</b>	MCAM antibody was produced in Mouse
<b>Clonality:</b>	Monoclonal
<b>Isotype:</b>	IgG1
<b>Conjugations:</b>	Fluorescein (FITC)
<b>Immunogen Species:</b>	MCAM / CD146 antibody was raised against Human
<b>Antigen Type:</b>	Recombinant protein
<b>Immunogen:</b>	MCAM / CD146 antibody was raised against recombinant human MUC18 (D1-D5) Fc protein.
<b>Specificity:</b>	Recognizes the 118kD cell surface glycoprotein CD146, also known as MUC18, Mel-CAM and S-endo.
<b>Reactivity:</b>	Human
<b>Purification:</b>	Protein G purified
<b>Presentation:</b>	PBS, pH 7.2, 1% BSA, 0.1% sodium azide.
<b>Recommended Storage:</b>	Long term: -20°C; Short term: +4°C. Avoid repeat freeze-thaw cycles.
<b>Usage Summary:</b>	Suitable for use in Flow Cytometry. Flow Cytometry: Neat-1:10. Labels 1E6 cells in 100ul.
<b>Uses:</b>	Flow Cytometry (Optimal dilution to be determined by the researcher)
<b>Size:</b>	100 µg
<b>Requested From:</b>	Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/25/2014

© 2014 LifeSpan BioSciences