


NDRG1 Rabbit anti-Human Polyclonal (aa19-48) Antibody - LS-C164789 - LSBio


CatalogID:	LS-C164789
Target:	N-myc downstream regulated 1 (NDRG1)
Synonyms:	NDRG1 Antibody, CMT4D Antibody, DRG-1 Antibody, GC4 Antibody, HMSNL Antibody, Protein regulated by oxygen-1 Antibody, PROXY1 Antibody, Protein NDRG1 Antibody, TARG1 Antibody, TDD5 Antibody, NDR1 Antibody, RTP Antibody, CAP43 Antibody, N-myc downstream regulated 1 Antibody, NMSL Antibody, RIT42 Antibody
Host	NDRG1 antibody was produced in Rabbit
Clonality:	Polyclonal
Immunogen Species:	NDRG1 antibody was raised against Human
Antigen Type:	Synthetic peptide
Immunogen:	NDRG1 antibody was raised against kLH-conjugated synthetic peptide from N-terminal region of human NDRG1.
Specificity:	Human NDRG1
Epitope:	aa19-48
Reactivity:	Human
Purification:	Immunoaffinity purified
Presentation:	PBS, 0.09% sodium azide
Recommended Storage:	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.
Uses:	IHC - Paraffin (1:50 - 1:100), Western blot (1:1000), Flow Cytometry (1:10 - 1:50) (Optimal dilution to be determined by the researcher)
Size:	400 µl

Immunohistochemistry Image:




Formalin-fixed and paraffin-embedded human prostate carcinoma with NDRG1 Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining.

Flow Cytometry Image:



Flow cytometry of MDA-231 cells using NDRG1 Antibody (N-term)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Western Blot Image:



Western blot of NDRG1 Antibody (N-term) in MDA-MB231 cell line lysates (35 ug/lane). NDRG1 (arrow) was detected using the purified Pab.

Requested From:

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/26/2014

© 2014 LifeSpan BioSciences